

ELLIPTIC DEFORMED SUPERALGEBRA $U_{q,p}(\widehat{sl}(M|N))$

January 18, 2013

TAKEO KOJIMA

*Department of Mathematics and Physics, Graduate School of Science and Engineering,
Yamagata University, Jonan 4-3-16, Yonezawa 992-8510, Japan
kojima@yz.yamagata-u.ac.jp*

Abstract

We introduce the elliptic superalgebra $U_{q,p}(\widehat{sl}(M|N))$ as one parameter deformation of the quantum superalgebra $U_q(\widehat{sl}(M|N))$. For an arbitrary level $k \neq 1$ we give the bosonization of the elliptic superalgebra $U_{q,p}(\widehat{sl}(1|2))$ and the screening currents that commute with $U_{q,p}(\widehat{sl}(1|2))$ modulo total difference.

1 Introduction

Infinite dimensional symmetry has been an impressive success in conformal field theory (CFT) [1]. Solvable lattice model is an off-critical extension of CFT and infinite dimensional symmetry plays an important role in algebraic analysis of solvable lattice model [2]. The lattice counterpart of minimal unitary CFT is Andrews-Baxter-Forrester (ABF) model [3], whose Boltzmann weights are elliptic solutions of the Yang-Baxter equation (YBE) of the face-type. Among the solvable models based on YBE, those related to elliptic solutions occupy a fundamental place.

Elliptic algebras are certain algebraic structures introduced to investigate these elliptic models. In study of k -fusion hierarchy of ABF model, Konno [4] introduced the elliptic algebra $U_{q,p}(\widehat{sl}(2))$ and constructed bosonization of the vertex operator by using this algebra. Jimbo-Konno-Odake-Shiraishi [5] constructed the elliptic algebra $U_{q,p}(g)$ by dressing the usual Drinfeld currents [20] of the quantum group $U_q(g)$ for non-twisted affine Lie algebra g . In this paper we introduce the elliptic deformed superalgebra $U_{q,p}(\widehat{sl}(M|N))$ as one parameter deformation of the quantum superalgebra $U_q(\widehat{sl}(M|N))$. We give the bosonization of the elliptic superalgebra $U_{q,p}(\widehat{sl}(1|2))$ and $U_{q,p}(\widehat{sl}(2|1))$ for generic level k , and give the screening currents that commute with $U_{q,p}(\widehat{sl}(1|2))$ and $U_{q,p}(\widehat{sl}(2|1))$ modulo total difference.

In this paper we aim to contribute mathematical tools for the study of super $\widehat{sl}(M|N)$ -family of the ABF model [19]. Mathematical tools are the elliptic algebra $U_{q,p}(\widehat{sl}(M|N))$ and bosonizations. We give comments on $\widehat{sl}(N)$ -family of the ABF model, where such mathematical tools have been used previously in analogous, but simpler case. Andrews-Baxter-Forrester [3] introduced the ABF model, that gives an extension of the hard hexagon model, and derived local height probabilities by Baxter's corner transfer matrix method (CTM) [6]. The k -fusion and higher-rank generalization, that we call $\widehat{sl}(N)$ -family of the ABF model, have been studied in [7, 8, 10, 11]. Inspired by the vertex operator approach to the 6-vertex model [2, 12, 13, 14], that originated from CTM, Lukyanov-Pugai [15] studied the vertex operator approach to the ABF model, and derived integral representations of multi-point local height probabilities. In study of k -fusion hierarchy of ABF model, Konno [4] introduced the elliptic algebra $U_{q,p}(\widehat{sl}(2))$ and constructed bosonization of the vertex operator by using this algebra. The vertex operator approach to the higher-rank generalization of the ABF model have been studied in [16, 17, 18]. In the vertex operator approach to $\widehat{sl}(N)$ -family of the ABF model, bosonization of the vertex operator played important role. In construction of the vertex operator, the current of the elliptic algebra $U_{q,p}(\widehat{sl}(N))$ and its bosonization played important roles. In order to derive integral representation of multi-point local height probabilities of super $\widehat{sl}(M|N)$ -family of the ABF model, we have to construct bosonizations of the vertex operators by using the current of the elliptic algebra $U_{q,p}(\widehat{sl}(M|N))$, and understand the structure of the space of state of the model by CTM [6], that has been open problem for superalgebra $\widehat{sl}(M|N)$.

Next we give comments on pure mathematical aspects. Through an attempt to understand solvable models based on elliptic solutions of YBE, various versions of elliptic algebras have been introduced [4, 5, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30]. It is important to understand not only themselves but also relations between them. Here we summarize some basic facts on the elliptic quantum group $\mathcal{B}_{q,\lambda}(g)$ and the elliptic algebra $U_{q,p}(g)$. The elliptic quantum group

$\mathcal{B}_{q,\lambda}(g)$, was introduced by twisting the standard quantum group $U_q(g)$ [25, 26, 27, 28, 29], where g is the symmetrizable Kac-Moody algebra. The elliptic quantum group $\mathcal{B}_{q,\lambda}(g)$ has quasi-Hopf structure and the elliptic algebra $U_{q,p}(\widehat{sl}(2))$ has H -Hopf algebroid structure [31, 32]. The realizations of the L -operators of the elliptic quantum group $\mathcal{B}_{q,\lambda}(g)$ were constructed in [5, 21, 22] by using the currents of the elliptic algebra $U_{q,p}(g)$ for $g = \widehat{sl}(N), A_2^{(2)}$. This suggests that the currents of $U_{q,p}(g)$ give the Drinfeld currents [20] of the elliptic quantum group $\mathcal{B}_{q,\lambda}(g)$. The construction of the elliptic quantum group $\mathcal{B}_{q,\lambda}(g)$ has been extended to the superalgebra $g = \widehat{sl}(M|N)$ [30]. In this paper we introduce the elliptic algebra $U_{q,p}(\widehat{sl}(M|N))$. We conjecture that the L -operator of $\mathcal{B}_{q,\lambda}(\widehat{sl}(M|N))$ is constructed by using the currents of $U_{q,p}(\widehat{sl}(M|N))$ and that there exists H -Hopf algebroid structure for $U_{q,p}(\widehat{sl}(M|N))$. The bosonizations of the vertex operators give useful information for construction of the L -operator, thorough so-called Miki's construction [33] of the L -operator. The above is background mathematical theory of the vertex operator. Next we give a comment on mathematical phenomenon of the space of state. Date-Jimbo-Kuniba-Miwa-Okado [8, 9, 10, 11] found that local height probabilities of $\widehat{sl}(N)$ -family of the ABF model were expressed in terms of the branching coefficients appearing in the irreducible decomposition of the character of $\widehat{sl}(N)$ [34, 35]. In order to extend this to super $\widehat{sl}(M|N)$ -family of the ABF model, we have to know the character formulae of the superalgebra $\widehat{sl}(M|N)$, that gives the affine generalization of formulae [36, 37].

The text is organized as follows. In section 2, after preparing the notations and giving the definition of the quantum group $U_q(\widehat{sl}(M|N))$, we introduce the elliptic deformed superalgebra $U_{q,p}(\widehat{sl}(M|N))$. Our approach is based on the dressing procedure of the Drinfeld current of the quantum group. In section 3 we give bosonizations of the superalgebra $U_q(g), U_{q,p}(g)$ ($g = \widehat{sl}(1|2), \widehat{sl}(2|1)$) for an arbitrary level k . We give the screening currents that commute with $U_q(g), U_{q,p}(g)$ ($g = \widehat{sl}(1|2), \widehat{sl}(2|1)$) modulo total difference. In appendix we summarize some useful formulae of bosonizations and screening currents.

2 Elliptic deformed superalgebra $U_{q,p}(\widehat{sl}(M|N))$

In this section we introduce the elliptic superalgebra $U_{q,p}(\widehat{sl}(M|N))$. Kac [38] introduced the superalgebra generalization of contragredient Lie algebra. Van de Leur [39] classified the contragredient superalgebra g of finite growth. Yamane [40] introduced quantum affine superalgebra $U_q(g)$ and constructed the Drinfeld currents. We give elliptic deformation of the quantum affine superalgebra by developing the dressing procedure [5].

2.1 Quantum superalgebra $U_q(\widehat{sl}(M|N))$

In this section we review the Drinfeld realization of the quantum superalgebra $U_q(\widehat{sl}(M|N))$ for $M, N = 1, 2, 3, \dots$ [40]. We restrict our consideration to $M \neq N$. The quantum superalgebra $U_q(\widehat{sl}(M|N))$ in [40] is a q -deformation of the universal enveloping algebra of $\widehat{sl}(M|N)$ [39]. Hereafter we fix a complex number $q \neq 0, |q| < 1$. Let us set

$$[x, y] = xy - yx, \quad \{x, y\} = xy + yx, \quad [a]_q = \frac{q^a - q^{-a}}{q - q^{-1}}. \quad (2.1)$$

The Cartan matrix of the Lie superalgebra $\widehat{sl}(M|N)$ is given by

$$(A_{i,j})_{0 \leq i,j \leq M+N-1} = \begin{pmatrix} 0 & -1 & 0 & \cdots & & & \cdots & 0 & 1 \\ -1 & 2 & -1 & \cdots & & & \cdots & 0 & 0 \\ 0 & -1 & 2 & \cdots & & & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots & -1 & \cdots & & & \\ & & \cdots & -1 & 2 & -1 & \cdots & & \\ & & & \cdots & -1 & 0 & 1 & \cdots & \\ & & & & \cdots & 1 & -2 & 1 & \cdots \\ & & & & & \cdots & 1 & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & & & & \cdots & -2 & 1 & 0 \\ 0 & 0 & \cdots & & & & \cdots & 1 & -2 & 1 \\ 1 & 0 & \cdots & & & & \cdots & 0 & 1 & -2 \end{pmatrix}, \quad (2.2)$$

where the diagonal part is $(A_{i,i})_{0 \leq i \leq M+N-1} = (0, \overbrace{2, \dots, 2}^{M-1}, 0, \overbrace{-2, \dots, -2}^{N-1})$.

Definition 2.1 [40] *The generators of the quantum superalgebra $U_q(\widehat{sl}(M|N))$, which we call the Drinfeld generators, are given by*

$$x_{i,m}^\pm, a_{i,n}, h_i, c, \quad (1 \leq i \leq M+N-1, m \in \mathbb{Z}, n \in \mathbb{Z}_{\neq 0}). \quad (2.3)$$

Defining relations are

$$c : \text{central}, [h_i, a_{j,m}] = 0, \quad (2.4)$$

$$[a_{i,m}, a_{j,n}] = \frac{[A_{i,j}m]_q [cm]_q}{m} q^{-c|m|} \delta_{m+n,0}, \quad (2.5)$$

$$[h_i, x_j^\pm(z)] = \pm A_{i,j} x_j^\pm(z), \quad (2.6)$$

$$[a_{i,m}, x_j^+(z)] = \frac{[A_{i,j}m]_q}{m} q^{-c|m|} z^m x_j^+(z), \quad (2.7)$$

$$[a_{i,m}, x_j^-(z)] = -\frac{[A_{i,j}m]_q}{m} z^m x_j^-(z), \quad (2.8)$$

$$(z_1 - q^{\pm A_{i,j}} z_2) x_i^{\pm}(z_1) x_j^{\pm}(z_2) = (q^{\pm A_{j,i}} z_1 - z_2) x_j^{\pm}(z_2) x_i^{\pm}(z_1), \quad \text{for } |A_{i,j}| \neq 0, \quad (2.9)$$

$$x_i^{\pm}(z_1) x_j^{\pm}(z_2) = x_j^{\pm}(z_2) x_i^{\pm}(z_1), \quad \text{for } |A_{i,j}| = 0, (i, j) \neq (M, M), \quad (2.10)$$

$$\{x_M^{\pm}(z_1), x_M^{\pm}(z_2)\} = 0, \quad (2.11)$$

$$[x_i^+(z_1), x_j^-(z_2)] = \frac{\delta_{i,j}}{(q - q^{-1})z_1 z_2} \left(\delta(q^{-c} z_1/z_2) \psi_i^+(q^{\frac{c}{2}} z_2) - \delta(q^c z_1/z_2) \psi_i^-(q^{-\frac{c}{2}} z_2) \right),$$

for $(i, j) \neq (M, M),$ (2.12)

$$\{x_M^+(z_1), x_M^-(z_2)\} = \frac{1}{(q - q^{-1})z_1 z_2} \left(\delta(q^{-c} z_1/z_2) \psi_M^+(q^{\frac{c}{2}} z_2) - \delta(q^c z_1/z_2) \psi_M^-(q^{-\frac{c}{2}} z_2) \right), \quad (2.13)$$

$$\begin{aligned} & \left(x_i^{\pm}(z_1) x_i^{\pm}(z_2) x_j^{\pm}(z) - (q + q^{-1}) x_i^{\pm}(z_1) x_j^{\pm}(z) x_i^{\pm}(z_2) + x_j^{\pm}(z) x_i^{\pm}(z_1) x_i^{\pm}(z_2) \right) \\ & + (z_1 \leftrightarrow z_2) = 0, \quad \text{for } |A_{i,j}| = 1, i \neq M, \end{aligned} \quad (2.14)$$

$$\begin{aligned} & (x_M^{\pm}(z_1) x_{M+1}^{\pm}(w_1) x_M^{\pm}(z_2) x_{M-1}^{\pm}(w_2) - q^{-1} x_M^{\pm}(z_1) x_{M+1}^{\pm}(w_1) x_{M-1}^{\pm}(w_2) x_M^{\pm}(z_2) \\ & - q x_M^{\pm}(z_1) x_M^{\pm}(z_2) x_{M-1}^{\pm}(w_2) x_{M+1}^{\pm}(w_1) + x_M^{\pm}(z_1) x_{M-1}^{\pm}(w_2) x_M^{\pm}(z_2) x_{M+1}^{\pm}(w_1) \\ & + x_{M+1}^{\pm}(w_1) x_M^{\pm}(z_2) x_{M-1}^{\pm}(w_2) x_M^{\pm}(z_1) - q^{-1} x_{M+1}^{\pm}(w_1) x_{M-1}^{\pm}(w_2) x_M^{\pm}(z_2) x_M^{\pm}(z_1) \\ & - q x_M^{\pm}(z_2) x_{M-1}^{\pm}(w_2) x_{M+1}^{\pm}(w_1) x_M^{\pm}(z_1) + x_{M-1}^{\pm}(w_2) x_M^{\pm}(z_2) x_{M+1}^{\pm}(w_1) x_M^{\pm}(z_1)) \\ & + (z_1 \leftrightarrow z_2) = 0, \end{aligned} \quad (2.15)$$

where we have used $\delta(z) = \sum_{m \in \mathbb{Z}} z^m$. Here we have set the generating functions

$$x_j^{\pm}(z) = \sum_{m \in \mathbb{Z}} x_{j,m}^{\pm} z^{-m-1}, \quad (2.16)$$

$$\psi_i^+(q^{\frac{c}{2}} z) = q^{h_i} \exp \left((q - q^{-1}) \sum_{m > 0} a_{i,m} z^{-m} \right), \quad (2.17)$$

$$\psi_i^-(q^{-\frac{c}{2}} z) = q^{-h_i} \exp \left(-(q - q^{-1}) \sum_{m > 0} a_{i,-m} z^m \right). \quad (2.18)$$

We changed the gauge of boson $a_{i,m}$ from those of [40]. In what follows we assume $c \in \mathbb{C}$.

2.2 Elliptic deformed superalgebra $U_{q,p}(\widehat{sl}(M|N))$

In this section we introduce the elliptic superalgebra $U_{q,p}(\widehat{sl}(M|N))$ for $M, N = 1, 2, 3, \dots, (M \neq N)$. Let us introduce a deformation parameter r such that

$$r, r^* = r - c > 0. \quad (2.19)$$

We often use the parameterization.

$$p = q^{2r} = e^{-\frac{2\pi i}{\tau}}, p^* = q^{2r^*} = e^{-\frac{2\pi i}{\tau^*}}, z = q^{2u}, w = q^{2v}. \quad (2.20)$$

We have $r\tau = r^*\tau^*$. Let us set the Jacobi theta functions $[u]$, $[u]^*$ by

$$[u] = q^{\frac{u^2}{r}-u} \frac{\Theta_p(q^{2u})}{(p;p)_\infty^3}, \quad [u]^* = q^{\frac{u^2}{r^*}-u} \frac{\Theta_{p^*}(q^{2u})}{(p^*;p^*)_\infty^3}. \quad (2.21)$$

Here we have used the standard symbols.

$$\Theta_p(z) = (p;p)_\infty (z;p)_\infty (pz^{-1};p)_\infty, \quad (2.22)$$

$$(z; t_1, \dots, t_k)_\infty = \prod_{n_1, \dots, n_k \geq 0} (1 - z t_1^{n_1} \dots t_k^{n_k}). \quad (2.23)$$

Definition 2.2 *The elliptic superalgebra $U_{q,p}(\widehat{sl}(M|N))$ is generated by the currents (operator valued function) and elements*

$$E_j(z), F_j(z), B_{j,n}, h_j, c \quad (1 \leq j \leq M+N-1, n \in \mathbb{Z}_{\neq 0}). \quad (2.24)$$

The defining relations are given as follows.

For $1 \leq i, j \leq M+N-1$, the relations are

$$c : \text{central}, [h_i, B_{j,m}] = 0, \quad (2.25)$$

$$[B_{i,m}, B_{j,n}] = \frac{[A_{i,j}m]_q [cm]_q}{m} \frac{[r^*m]_q}{[rm]_q} \delta_{m+n,0}, \quad (2.26)$$

$$[h_i, E_j(z)] = A_{i,j} E_j(z), [h_i, F_j(z)] = -A_{i,j} F_j(z), \quad (2.27)$$

$$[B_{i,m}, E_j(z)] = \frac{[A_{i,j}m]_q}{m} z^m E_j(z), [B_{i,m}, F_j(z)] = -\frac{[A_{i,j}m]_q}{m} \frac{[r^*m]_q}{[rm]_q} z^m F_j(z). \quad (2.28)$$

For $1 \leq i, j \leq M+N-1$ such that $(i, j) \neq (M, M)$, the relations are

$$\left[u_1 - u_2 - \frac{A_{i,j}}{2} \right]^* E_i(z_1) E_j(z_2) = \left[u_1 - u_2 + \frac{A_{i,j}}{2} \right]^* E_j(z_2) E_i(z_1), \quad (2.29)$$

$$\left[u_1 - u_2 + \frac{A_{i,j}}{2} \right] F_i(z_1) F_j(z_2) = \left[u_1 - u_2 - \frac{A_{i,j}}{2} \right] F_j(z_2) F_i(z_1), \quad (2.30)$$

$$[E_i(z_1), F_j(z_2)] = \frac{\delta_{i,j}}{(q - q^{-1})z_1 z_2} (\delta(q^{-c} z_1/z_2) H_i(q^r z_2) - \delta(q^c z_1/z_2) H_i(q^{-r} z_2)), \quad (2.31)$$

$$\{E_M(z_1), E_M(z_2)\} = 0, \quad \{F_M(z_1), F_M(z_2)\} = 0, \quad (2.32)$$

$$\{E_M(z_1), F_M(z_2)\} = \frac{1}{(q - q^{-1})z_1 z_2} (\delta(q^{-c} z_1/z_2) H_M(q^r z_2) - \delta(q^c z_1/z_2) H_M(q^{-r} z_2)). \quad (2.33)$$

For $1 \leq i, j \leq M+N-1$, the relations are

$$H_i(z_1) H_j(z_2) = \frac{[u_2 - u_1 - \frac{A_{i,j}}{2}]^* [u_2 - u_1 + \frac{A_{i,j}}{2}]}{[u_2 - u_1 + \frac{A_{i,j}}{2}]^* [u_2 - u_1 - \frac{A_{i,j}}{2}]} H_j(z_2) H_i(z_1), \quad (2.34)$$

$$H_i(z_1)E_j(z_2) = \frac{[u_1 - u_2 + \frac{r^*}{2} + \frac{A_{i,j}}{2}]^*}{[u_1 - u_2 + \frac{r^*}{2} - \frac{A_{i,j}}{2}]^*} E_j(z_2)H_i(z_1), \quad (2.35)$$

$$H_i(z_1)F_j(z_2) = \frac{[u_1 - u_2 + \frac{r}{2} + \frac{A_{i,j}}{2}]}{[u_1 - u_2 + \frac{r}{2} - \frac{A_{i,j}}{2}]} F_j(z_2)H_i(z_1). \quad (2.36)$$

For $1 \leq i, j \leq M + N - 1$ ($i \neq M$) such that $|A_{i,j}| = 1$, they satisfy the Serre relations

$$\begin{aligned} & \left(E_i(z_1)E_i(z_2)E_j(z) \frac{\{q^{A_{i,j}} \frac{z}{z_1}\}^* \{q^{A_{i,j}} \frac{z}{z_2}\}^*}{\{q^{-A_{i,j}} \frac{z}{z_1}\}^* \{q^{-A_{i,j}} \frac{z}{z_2}\}^*} \left(\frac{z}{z_2} \right)^{\frac{1}{r^*} A_{i,j}} \right. \\ & - (q + q^{-1})E_i(z_1)E_j(z)E_i(z_2) \frac{\{q^{A_{i,j}} \frac{z}{z_1}\}^* \{q^{A_{i,j}} \frac{z_2}{z}\}^*}{\{q^{-A_{i,j}} \frac{z}{z_1}\}^* \{q^{-A_{i,j}} \frac{z_2}{z}\}^*} \\ & + E_j(z)E_i(z_1)E_i(z_2) \frac{\{q^{A_{i,j}} \frac{z_1}{z}\}^* \{q^{A_{i,j}} \frac{z_2}{z}\}^*}{\{q^{-A_{i,j}} \frac{z_1}{z}\}^* \{q^{-A_{i,j}} \frac{z_2}{z}\}^*} \left(\frac{z_1}{z} \right)^{\frac{1}{r^*} A_{i,j}} \left. \right) \frac{\{q^{A_{i,i}} \frac{z_2}{z_1}\}^*}{\{q^{-A_{i,i}} \frac{z_2}{z_1}\}^*} z_1^{-\frac{1}{r^*} (A_{i,i} + A_{i,j})} \\ & + (z_1 \leftrightarrow z_2) = 0, \end{aligned} \quad (2.37)$$

$$\begin{aligned} & \left(F_i(z_1)F_i(z_2)F_j(z) \frac{\{q^{-A_{i,j}} \frac{z}{z_1}\} \{q^{-A_{i,j}} \frac{z}{z_2}\}}{\{q^{A_{i,j}} \frac{z}{z_1}\} \{q^{A_{i,j}} \frac{z}{z_2}\}} \left(\frac{z_2}{z} \right)^{\frac{1}{r}} \right. \\ & - (q + q^{-1})F_i(z_1)F_j(z)F_i(z_2) \frac{\{q^{-A_{i,j}} \frac{z}{z_1}\} \{q^{-A_{i,j}} \frac{z_2}{z}\}}{\{q^{A_{i,j}} \frac{z}{z_1}\} \{q^{A_{i,j}} \frac{z_2}{z}\}} \\ & + F_j(z)F_i(z_1)F_i(z_2) \frac{\{q^{-A_{i,j}} \frac{z_1}{z}\} \{q^{-A_{i,j}} \frac{z_2}{z}\}}{\{q^{A_{i,j}} \frac{z_1}{z}\} \{q^{A_{i,j}} \frac{z_2}{z}\}} \left(\frac{z}{z_1} \right)^{\frac{1}{r}} \left. \right) \frac{\{q^{-A_{i,i}} \frac{z_2}{z_1}\}}{\{q^{A_{i,i}} \frac{z_2}{z_1}\}} z_1^{\frac{1}{r} (A_{i,i} + A_{i,j})} \\ & + (z_1 \leftrightarrow z_2) = 0, \end{aligned} \quad (2.38)$$

and

$$\begin{aligned} & \left(E_M(z_1)E_{M+1}(w_1)E_M(z_2)E_{M-1}(w_2) \frac{\{\frac{qw_1}{z_1}\}^* \{\frac{qz_2}{w_1}\}^* \{\frac{w_2}{qz_1}\}^* \{\frac{w_2}{qz_2}\}^*}{\{\frac{w_1}{qz_1}\}^* \{\frac{z_2}{qw_1}\}^* \{\frac{qw_2}{z_1}\}^* \{\frac{qw_2}{z_2}\}^*} \left(\frac{w_2}{z_2} \right)^{\frac{1}{r^*}} \right. \\ & - q^{-1}E_M(z_1)E_{M+1}(w_1)E_{M-1}(w_2)E_M(z_2) \frac{\{\frac{qw_1}{z_1}\}^* \{\frac{w_2}{qz_1}\}^* \{\frac{qz_2}{w_1}\}^* \{\frac{z_2}{qw_2}\}^*}{\{\frac{w_1}{qz_1}\}^* \{\frac{qw_2}{z_1}\}^* \{\frac{z_2}{qw_1}\}^* \{\frac{qz_2}{w_2}\}^*} \\ & - qE_M(z_1)E_M(z_2)E_{M-1}(w_2)E_{M+1}(w_1) \frac{\{\frac{w_2}{qz_1}\}^* \{\frac{w_2}{qz_2}\}^* \{\frac{qw_1}{z_1}\}^* \{\frac{qw_1}{z_2}\}^*}{\{\frac{qw_2}{z_2}\}^* \{\frac{qw_1}{z_1}\}^* \{\frac{w_1}{qw_2}\}^* \{\frac{z_2}{qw_2}\}^*} \left(\frac{w_2}{w_1} \right)^{\frac{1}{r^*}} \\ & + E_M(z_1)E_{M-1}(w_2)E_M(z_2)E_{M+1}(w_1) \frac{\{\frac{w_2}{qz_1}\}^* \{\frac{z_2}{qw_1}\}^* \{\frac{qw_1}{z_1}\}^* \{\frac{qw_1}{z_2}\}^*}{\{\frac{qw_2}{z_1}\}^* \{\frac{qz_2}{z_1}\}^* \{\frac{w_1}{qw_1}\}^* \{\frac{w_1}{qw_2}\}^*} \left(\frac{z_2}{w_1} \right)^{\frac{1}{r^*}} \\ & + E_{M+1}(w_1)E_M(z_2)E_{M-1}(w_2)E_M(z_1) \frac{\{\frac{qz_2}{w_1}\}^* \{\frac{w_2}{qz_2}\}^* \{\frac{qz_1}{w_1}\}^* \{\frac{z_1}{qw_2}\}^*}{\{\frac{z_2}{qw_1}\}^* \{\frac{qw_2}{z_2}\}^* \{\frac{z_1}{qw_1}\}^* \{\frac{qz_1}{w_2}\}^*} \left(\frac{w_1}{z_2} \right)^{\frac{1}{r^*}} \\ & - q^{-1}E_{M+1}(w_1)E_{M-1}(w_2)E_M(z_2)E_M(z_1) \frac{\{\frac{qz_2}{w_1}\}^* \{\frac{z_2}{qw_2}\}^* \{\frac{qz_1}{w_1}\}^* \{\frac{z_1}{qw_2}\}^*}{\{\frac{z_2}{qw_1}\}^* \{\frac{qz_2}{w_2}\}^* \{\frac{z_1}{qw_1}\}^* \{\frac{qz_1}{w_2}\}^*} \left(\frac{w_1}{w_2} \right)^{\frac{1}{r^*}} \\ & - qE_M(z_2)E_{M-1}(w_2)E_{M+1}(w_1)E_M(z_1) \frac{\{\frac{w_2}{qz_2}\}^* \{\frac{w_1}{qw_2}\}^* \{\frac{qz_1}{w_2}\}^* \{\frac{qz_1}{w_1}\}^*}{\{\frac{qw_2}{z_2}\}^* \{\frac{qw_1}{z_2}\}^* \{\frac{z_1}{qw_2}\}^* \{\frac{z_1}{qw_1}\}^*} \\ & + E_{M-1}(w_2)E_M(z_2)E_{M+1}(w_1)E_M(z_1) \frac{\{\frac{z_2}{qw_2}\}^* \{\frac{qw_1}{z_1}\}^* \{\frac{z_1}{qw_2}\}^* \{\frac{qz_1}{w_1}\}^*}{\{\frac{qz_2}{w_2}\}^* \{\frac{w_1}{qw_1}\}^* \{\frac{qz_1}{w_2}\}^* \{\frac{z_1}{qw_1}\}^*} \left(\frac{z_2}{w_2} \right)^{\frac{1}{r^*}} \left. \right) \end{aligned}$$

$$+(z_1 \leftrightarrow z_2) = 0, \quad (2.39)$$

$$\begin{aligned}
& \left(F_M(z_1)F_{M+1}(w_1)F_M(z_2)F_{M-1}(w_2) \frac{\left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{z_2}{qw_1} \right\} \left\{ \frac{qw_2}{z_1} \right\} \left\{ \frac{qw_2}{z_2} \right\}}{\left\{ \frac{qw_1}{z_1} \right\} \left\{ \frac{qz_2}{w_1} \right\} \left\{ \frac{w_2}{qz_1} \right\} \left\{ \frac{w_2}{qw_2} \right\}} \left(\frac{z_2}{w_2} \right)^{\frac{1}{r}} \right. \\
& - q^{-1} F_M(z_1)F_{M+1}(w_1)F_{M-1}(w_2)F_M(z_2) \frac{\left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{qw_2}{z_1} \right\} \left\{ \frac{z_2}{qw_1} \right\} \left\{ \frac{qz_2}{w_2} \right\}}{\left\{ \frac{qw_1}{z_1} \right\} \left\{ \frac{w_2}{qz_1} \right\} \left\{ \frac{qz_2}{w_1} \right\} \left\{ \frac{z_2}{qw_2} \right\}} \\
& - q F_M(z_1)F_M(z_2)F_{M-1}(w_2)F_{M+1}(w_1) \frac{\left\{ \frac{qw_2}{z_1} \right\} \left\{ \frac{qw_2}{z_2} \right\} \left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{w_1}{qw_2} \right\}}{\left\{ \frac{w_2}{qz_2} \right\} \left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{qw_1}{z_2} \right\} \left\{ \frac{qz_2}{w_2} \right\}} \left(\frac{w_1}{w_2} \right)^{\frac{1}{r}} \\
& + F_M(z_1)F_{M-1}(w_2)F_M(z_2)F_{M+1}(w_1) \frac{\left\{ \frac{qw_2}{z_1} \right\} \left\{ \frac{qz_2}{z_1} \right\} \left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{w_1}{qw_2} \right\}}{\left\{ \frac{w_2}{qz_1} \right\} \left\{ \frac{z_2}{qz_1} \right\} \left\{ \frac{qw_1}{z_1} \right\} \left\{ \frac{qw_1}{z_2} \right\}} \left(\frac{w_1}{z_2} \right)^{\frac{1}{r}} \\
& + F_{M+1}(w_1)F_M(z_2)F_{M-1}(w_2)F_M(z_1) \frac{\left\{ \frac{z_2}{qw_1} \right\} \left\{ \frac{qw_2}{z_2} \right\} \left\{ \frac{z_1}{qw_1} \right\} \left\{ \frac{qz_1}{w_2} \right\}}{\left\{ \frac{qz_2}{w_1} \right\} \left\{ \frac{w_2}{qz_2} \right\} \left\{ \frac{qz_1}{w_1} \right\} \left\{ \frac{z_1}{qw_2} \right\}} \left(\frac{z_2}{w_1} \right)^{\frac{1}{r}} \\
& - q^{-1} F_{M+1}(w_1)F_{M-1}(w_2)F_M(z_2)F_M(z_1) \frac{\left\{ \frac{z_2}{qw_1} \right\} \left\{ \frac{qz_2}{w_2} \right\} \left\{ \frac{z_1}{qw_1} \right\} \left\{ \frac{qz_1}{w_2} \right\}}{\left\{ \frac{qz_2}{w_1} \right\} \left\{ \frac{z_2}{qw_2} \right\} \left\{ \frac{qz_1}{w_1} \right\} \left\{ \frac{z_1}{qw_2} \right\}} \left(\frac{w_2}{w_1} \right)^{\frac{1}{r}} \\
& - q F_M(z_2)F_{M-1}(w_2)F_{M+1}(w_1)F_M(z_1) \frac{\left\{ \frac{qw_2}{z_2} \right\} \left\{ \frac{qw_1}{z_2} \right\} \left\{ \frac{z_1}{qw_2} \right\} \left\{ \frac{z_1}{qw_1} \right\}}{\left\{ \frac{w_2}{qz_2} \right\} \left\{ \frac{w_1}{qz_2} \right\} \left\{ \frac{qz_1}{w_2} \right\} \left\{ \frac{qz_1}{w_1} \right\}} \\
& + F_{M-1}(w_2)F_M(z_2)F_{M+1}(w_1)F_M(z_1) \frac{\left\{ \frac{qz_2}{w_2} \right\} \left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{qz_1}{w_2} \right\} \left\{ \frac{z_1}{qw_1} \right\}}{\left\{ \frac{z_2}{qw_2} \right\} \left\{ \frac{qw_1}{z_1} \right\} \left\{ \frac{z_1}{qw_2} \right\} \left\{ \frac{qz_1}{w_1} \right\}} \left(\frac{w_2}{z_2} \right)^{\frac{1}{r}} \Bigg) \\
& + (z_1 \leftrightarrow z_2) = 0. \quad (2.40)
\end{aligned}$$

Here we have used the abbreviations

$$\{z\}^* = (p^* z; p^*)_\infty, \quad \{z\} = (pz; p)_\infty. \quad (2.41)$$

2.3 Dressing construction

In this section we construct $U_{q,p}(\widehat{sl}(M|N))$ from $U_q(\widehat{sl}(M|N))$ by developing the dressing procedure [5].

Definition 2.3 Let us introduce the dressing operators $u_j^\pm(z, p)$, ($1 \leq j \leq M + N - 1$) by

$$u_j^+(z, p) = \exp \left(\sum_{m>0} \frac{1}{[r^* m]_q} a_{j,-m} (q^r z)^m \right), \quad (2.42)$$

$$u_j^-(z, p) = \exp \left(- \sum_{m>0} \frac{1}{[r m]_q} a_{j,m} (q^{-r} z)^{-m} \right). \quad (2.43)$$

Straightforward calculations show the following propositions.

Proposition 2.4 For $1 \leq i, j \leq M + N - 1$, we have

$$u_i^+(z_1, p)x_j^+(z_2) = \frac{(p^*q^{A_{i,j}}z_1/z_2 : p^*)_\infty}{(p^*q^{-A_{i,j}}z_1/z_2; p^*)_\infty} x_j^+(z_2)u_i^+(z_1, p), \quad (2.44)$$

$$u_i^+(z_1, p)x_j^-(z_2) = \frac{(p^*q^{-A_{i,j}+c}z_1/z_2 : p^*)_\infty}{(p^*q^{A_{i,j}+c}z_1/z_2; p^*)_\infty} x_j^-(z_2)u_i^+(z_1, p), \quad (2.45)$$

$$u_i^-(z_1, p)x_j^+(z_2) = \frac{(pq^{-A_{i,j}-c}z_1/z_2 : p)_\infty}{(pq^{A_{i,j}-c}z_1/z_2; p)_\infty} x_j^+(z_2)u_i^-(z_1, p), \quad (2.46)$$

$$u_i^-(z_1, p)x_j^-(z_2) = \frac{(pq^{A_{i,j}}z_1/z_2 : p)_\infty}{(pq^{-A_{i,j}}z_1/z_2; p)_\infty} x_j^-(z_2)u_i^-(z_1, p), \quad (2.47)$$

$$u_i^+(z_1, p)u_j^-(z_2, p) = \frac{(pq^{-A_{i,j}-c}z_1/z_2; p)_\infty (p^*q^{A_{i,j}+c}z_1/z_2; p^*)_\infty}{(pq^{A_{i,j}-c}z_1/z_2; p)_\infty (p^*q^{-A_{i,j}+c}z_1/z_2; p^*)_\infty} u_j^-(z_2, p)u_i^+(z_1, p). \quad (2.48)$$

Definition 2.5 We define the dressing currents $e_j(z, p), f_j(z, p), \psi_j^\pm(z, p)$, ($1 \leq j \leq M + N - 1$) by

$$e_j(z, p) = u_j^+(z, p)x_j^+(z), \quad (2.49)$$

$$f_j(z, p) = x_j^-(z)u_j^-(z, p), \quad (2.50)$$

$$\psi_j^+(z, p) = u_j^+(q^{\frac{c}{2}}z, p)\psi_j^+(z)u_j^-(q^{-\frac{c}{2}}z, p), \quad (2.51)$$

$$\psi_j^-(z, p) = u_j^+(q^{-\frac{c}{2}}z, p)\psi_j^-(z)u_j^-(q^{\frac{c}{2}}z, p). \quad (2.52)$$

Proposition 2.6 The currents $e_i(z, p), f_i(z, p)$ and $a_{i,n}, h_i, c$, ($1 \leq i \leq M + N - 1, n \in \mathbb{Z}_{\neq 0}$) satisfy the following relations

$$c : \text{central}, [h_i, a_{j,m}] = 0, \quad (2.53)$$

$$[a_{i,m}, a_{j,n}] = \frac{[A_{i,j}m]_q [cm]_q}{m} q^{-c|m|} \delta_{m+n,0}, \quad (2.54)$$

$$[h_i, e_j(z, p)] = A_{i,j}e_j(z, p), \quad [h_i, f_j(z, p)] = -A_{i,j}f_j(z, p), \quad (2.55)$$

$$[a_{i,m}, e_j(z, p)] = \frac{[A_{i,j}m]_q}{m} z^m e_j(z, p) \times \begin{cases} \frac{[rm]_q}{[r^*m]_q}, & (m > 0) \\ q^{cm}, & (m < 0) \end{cases}, \quad (2.56)$$

$$[a_{i,m}, f_j(z, p)] = -\frac{[A_{i,j}m]_q}{m} z^m f_j(z, p) \times \begin{cases} 1, & (m > 0) \\ \frac{[r^*m]_q}{[rm]_q} q^{cm}, & (m < 0) \end{cases}, \quad (2.57)$$

$$\begin{aligned} & z_1 \Theta_{p^*}(q^{A_{i,j}}z_2/z_1)e_i(z_1, p)e_j(z_2, p) \\ &= -z_2 \Theta_{p^*}(q^{A_{i,j}}z_2/z_1)e_j(z_2, p)e_i(z_1, p), \quad \text{for } |A_{i,j}| \neq 0, \end{aligned} \quad (2.58)$$

$$[e_i(z_1, p), e_j(z_2, p)] = 0, \quad \text{for } |A_{i,j}| = 0, (i, j) \neq (M, M), \quad (2.59)$$

$$\{e_M(z_1, p), e_M(z_2, p)\} = 0, \quad (2.60)$$

$$z_1 \Theta_p(q^{-A_{i,j}}z_2/z_1)f_i(z_1, p)f_j(z_2, p)$$

$$= -z_2 \Theta_p(q^{-A_{j,i}} z_2/z_1) f_j(z_2, p) f_i(z_1, p), \quad \text{for } |A_{i,j}| \neq 0, \quad (2.61)$$

$$[f_i(z_1, p), f_j(z_2, p)] = 0, \quad \text{for } |A_{i,j}| = 0, (i, j) \neq (M, M), \quad (2.62)$$

$$\{f_M(z_1, p), f_M(z_2, p)\} = 0, \quad (2.63)$$

$$[e_i(z_1, p), f_j(z_2, p)] = \frac{\delta_{i,j}}{(q - q^{-1}) z_1 z_2} \left(\delta(q^{-c} z_1/z_2) \psi_i^+(q^{\frac{c}{2}} z_2, p) - \delta(q^c z_1/z_2) \psi_i^-(q^{-\frac{c}{2}} z_2, p) \right),$$

for $(i, j) \neq (M, M),$ (2.64)

$$\{e_M(z_1, p), f_M(z_2, p)\} = \frac{1}{(q - q^{-1}) z_1 z_2} \left(\delta(q^{-c} z_1/z_2) \psi_M^+(q^{\frac{c}{2}} z_2, p) - \delta(q^c z_1/z_2) \psi_M^-(q^{-\frac{c}{2}} z_2, p) \right), \quad (2.65)$$

$$\begin{aligned} & \left(e_i(z_1, p) e_i(z_2, p) e_j(z, p) \frac{\{q^{A_{i,j}} \frac{z}{z_1}\}^* \{q^{A_{i,j}} \frac{z}{z_2}\}^*}{\{q^{-A_{i,j}} \frac{z}{z_1}\}^* \{q^{-A_{i,j}} \frac{z}{z_2}\}^*} \right. \\ & - (q + q^{-1}) e_i(z_1, p) e_j(z, p) e_i(z_2, p) \frac{\{q^{A_{i,j}} \frac{z}{z_1}\}^* \{q^{A_{i,j}} \frac{z_2}{z}\}^*}{\{q^{-A_{i,j}} \frac{z}{z_1}\}^* \{q^{-A_{i,j}} \frac{z_2}{z}\}^*} \\ & + e_j(z, p) e_i(z_1, p) e_i(z_2, p) \frac{\{q^{A_{i,j}} z_1/z\}^* \{q^{A_{i,j}} z_2/z\}^*}{\{q^{-A_{i,j}} \frac{z_1}{z}\}^* \{q^{-A_{i,j}} \frac{z_2}{z}\}^*} \left. \right) \frac{\{q^{A_{i,i}} \frac{z_2}{z_1}\}^*}{\{q^{-A_{i,i}} \frac{z_2}{z_1}\}^*} \\ & + (z_1 \leftrightarrow z_2) = 0, \quad \text{for } |A_{i,j}| = 1, i \neq M, \end{aligned} \quad (2.66)$$

$$\begin{aligned} & \left(f_i(z_1, p) f_i(z_2, p) f_j(z, p) \frac{\{q^{-A_{i,j}} \frac{z}{z_1}\} \{q^{-A_{i,j}} \frac{z}{z_2}\}}{\{q^{A_{i,j}} \frac{z}{z_1}\} \{q^{A_{i,j}} \frac{z}{z_2}\}} \right. \\ & - (q + q^{-1}) f_i(z_1, p) f_j(z, p) f_i(z_2, p) \frac{\{q^{-A_{i,j}} \frac{z}{z_1}\} \{q^{-A_{i,j}} \frac{z_2}{z}\}}{\{q^{A_{i,j}} \frac{z}{z_1}\} \{q^{A_{i,j}} \frac{z_2}{z}\}} \\ & + f_j(z, p) f_i(z_1, p) f_i(z_2, p) \frac{\{q^{-A_{i,j}} \frac{z_1}{z}\} \{q^{-A_{i,j}} \frac{z_2}{z}\}}{\{q^{A_{i,j}} \frac{z_1}{z}\} \{q^{A_{i,j}} \frac{z_2}{z}\}} \left. \right) \frac{\{q^{-A_{i,i}} \frac{z_2}{z_1}\}}{\{q^{A_{i,i}} \frac{z_2}{z_1}\}} \\ & + (z_1 \leftrightarrow z_2) = 0, \quad \text{for } |A_{i,j}| = 1, i \neq M, \end{aligned} \quad (2.67)$$

$$\begin{aligned} & \left(e_M(z_1, p) e_{M+1}(w_1, p) e_M(z_2, p) e_{M-1}(w_2, p) \frac{\{\frac{qw_1}{z_1}\}^* \{\frac{qz_2}{w_1}\}^* \{\frac{w_2}{qz_1}\}^* \{\frac{w_2}{qz_2}\}^*}{\{\frac{w_1}{qz_1}\}^* \{\frac{z_2}{qw_1}\}^* \{\frac{qw_2}{z_1}\}^* \{\frac{qw_2}{z_2}\}^*} \right. \\ & - q^{-1} e_M(z_1, p) e_{M+1}(w_1, p) e_{M-1}(w_2, p) e_M(z_2, p) \frac{\{\frac{qw_1}{z_1}\}^* \{\frac{w_2}{qz_1}\}^* \{\frac{qz_2}{w_1}\}^* \{\frac{z_2}{qw_2}\}^*}{\{\frac{w_1}{qz_1}\}^* \{\frac{qw_2}{z_1}\}^* \{\frac{z_2}{qw_1}\}^* \{\frac{qz_2}{w_2}\}^*} \\ & - q e_M(z_1, p) e_M(z_2, p) e_{M-1}(w_2, p) e_{M+1}(w_1, p) \frac{\{\frac{w_2}{qz_1}\}^* \{\frac{w_2}{qz_2}\}^* \{\frac{qw_1}{z_1}\}^* \{\frac{qw_1}{z_2}\}^*}{\{\frac{qw_2}{z_2}\}^* \{\frac{qw_1}{z_1}\}^* \{\frac{w_1}{qz_2}\}^* \{\frac{z_2}{qw_2}\}^*} \\ & + e_M(z_1, p) e_{M-1}(w_2, p) e_M(z_2, p) e_{M+1}(w_1, p) \frac{\{\frac{w_2}{qz_1}\}^* \{\frac{z_2}{qz_1}\}^* \{\frac{qw_1}{z_1}\}^* \{\frac{qw_1}{z_2}\}^*}{\{\frac{qw_2}{z_1}\}^* \{\frac{qz_2}{z_1}\}^* \{\frac{w_1}{qz_1}\}^* \{\frac{w_1}{qz_2}\}^*} \\ & + e_{M+1}(w_1, p) e_M(z_2, p) e_{M-1}(w_2, p) e_M(z_1, p) \frac{\{\frac{qz_2}{w_1}\}^* \{\frac{w_2}{qz_2}\}^* \{\frac{qz_1}{w_1}\}^* \{\frac{z_1}{qw_2}\}^*}{\{\frac{z_2}{qw_1}\}^* \{\frac{qw_2}{z_2}\}^* \{\frac{z_1}{qw_1}\}^* \{\frac{qz_1}{w_2}\}^*} \\ & - q^{-1} e_{M+1}(w_1, p) e_{M-1}(w_2, p) e_M(z_2, p) e_M(z_1, p) \frac{\{\frac{qz_2}{w_1}\}^* \{\frac{z_2}{qw_2}\}^* \{\frac{qz_1}{w_1}\}^* \{\frac{z_1}{qw_2}\}^*}{\{\frac{z_2}{qw_1}\}^* \{\frac{qz_2}{w_2}\}^* \{\frac{z_1}{qw_1}\}^* \{\frac{qz_1}{w_2}\}^*} \\ & - q e_M(z_2, p) e_{M-1}(w_2, p) e_{M+1}(w_1, p) e_M(z_1, p) \frac{\{\frac{w_2}{qz_2}\}^* \{\frac{w_1}{qz_2}\}^* \{\frac{qz_1}{w_2}\}^* \{\frac{qz_1}{w_1}\}^*}{\{\frac{qw_2}{z_2}\}^* \{\frac{qw_1}{z_2}\}^* \{\frac{z_1}{qw_2}\}^* \{\frac{z_1}{qw_1}\}^*} \end{aligned}$$

$$\begin{aligned}
& + e_{M-1}(w_2, p) e_M(z_2, p) e_{M+1}(w_1, p) e_M(z_1, p) \frac{\left\{ \frac{z_2}{qw_2} \right\}^* \left\{ \frac{qw_1}{z_1} \right\}^* \left\{ \frac{z_1}{qw_2} \right\}^* \left\{ \frac{qw_2}{w_1} \right\}^*}{\left\{ \frac{qz_2}{w_2} \right\}^* \left\{ \frac{w_1}{qz_1} \right\}^* \left\{ \frac{qz_1}{w_2} \right\}^* \left\{ \frac{z_1}{qw_1} \right\}^*} \\
& + (z_1 \leftrightarrow z_2) = 0,
\end{aligned} \tag{2.68}$$

$$\begin{aligned}
& \left(f_M(z_1, p) f_{M+1}(w_1, p) f_M(z_2, p) f_{M-1}(w_2, p) \frac{\left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{z_2}{qw_1} \right\} \left\{ \frac{qw_2}{z_1} \right\} \left\{ \frac{qw_2}{z_2} \right\}}{\left\{ \frac{qw_1}{z_1} \right\} \left\{ \frac{qz_2}{w_1} \right\} \left\{ \frac{w_2}{qz_1} \right\} \left\{ \frac{w_2}{qz_2} \right\}} \right. \\
& - q^{-1} f_M(z_1, p) f_{M+1}(w_1, p) f_{M-1}(w_2, p) f_M(z_2, p) \frac{\left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{qw_2}{z_1} \right\} \left\{ \frac{z_2}{qw_1} \right\} \left\{ \frac{qz_2}{w_2} \right\}}{\left\{ \frac{qw_1}{z_1} \right\} \left\{ \frac{w_2}{qz_1} \right\} \left\{ \frac{qz_2}{w_1} \right\} \left\{ \frac{z_2}{qw_2} \right\}} \\
& - q f_M(z_1, p) f_M(z_2, p) f_{M-1}(w_2, p) f_{M+1}(w_1, p) \frac{\left\{ \frac{qw_2}{z_1} \right\} \left\{ \frac{qw_2}{z_2} \right\} \left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{w_1}{qz_2} \right\}}{\left\{ \frac{w_2}{qz_2} \right\} \left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{qw_1}{z_2} \right\} \left\{ \frac{qw_2}{z_2} \right\}} \\
& + f_M(z_1, p) f_{M-1}(w_2, p) f_M(z_2, p) f_{M+1}(w_1, p) \frac{\left\{ \frac{qw_2}{z_1} \right\} \left\{ \frac{qz_2}{z_1} \right\} \left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{w_1}{qz_2} \right\}}{\left\{ \frac{w_2}{qz_1} \right\} \left\{ \frac{z_2}{qz_1} \right\} \left\{ \frac{qw_1}{z_1} \right\} \left\{ \frac{qw_1}{z_2} \right\}} \\
& + f_{M+1}(w_1, p) f_M(z_2, p) f_{M-1}(w_2, p) f_M(z_1, p) \frac{\left\{ \frac{z_2}{qw_1} \right\} \left\{ \frac{qw_2}{z_2} \right\} \left\{ \frac{z_1}{qw_1} \right\} \left\{ \frac{qz_1}{w_2} \right\}}{\left\{ \frac{qz_2}{w_1} \right\} \left\{ \frac{w_2}{qz_2} \right\} \left\{ \frac{qz_1}{w_1} \right\} \left\{ \frac{z_1}{qw_2} \right\}} \\
& - q^{-1} f_{M+1}(w_1, p) f_{M-1}(w_2, p) f_M(z_2, p) f_M(z_1, p) \frac{\left\{ \frac{z_2}{qw_1} \right\} \left\{ \frac{qz_2}{w_2} \right\} \left\{ \frac{z_1}{qw_1} \right\} \left\{ \frac{qz_1}{w_2} \right\}}{\left\{ \frac{qz_2}{w_1} \right\} \left\{ \frac{z_2}{qw_2} \right\} \left\{ \frac{qz_1}{w_1} \right\} \left\{ \frac{z_1}{qw_2} \right\}} \\
& - q f_M(z_2, p) f_{M-1}(w_2, p) f_{M+1}(w_1, p) f_M(z_1, p) \frac{\left\{ \frac{qw_2}{z_2} \right\} \left\{ \frac{qw_1}{z_2} \right\} \left\{ \frac{z_1}{qw_2} \right\} \left\{ \frac{z_1}{qw_1} \right\}}{\left\{ \frac{w_2}{qz_2} \right\} \left\{ \frac{w_1}{qz_2} \right\} \left\{ \frac{qz_1}{w_2} \right\} \left\{ \frac{qz_1}{w_1} \right\}} \\
& + f_{M-1}(w_2, p) f_M(z_2, p) f_{M+1}(w_1, p) f_M(z_1, p) \frac{\left\{ \frac{qz_2}{w_2} \right\} \left\{ \frac{w_1}{qz_1} \right\} \left\{ \frac{qz_1}{w_2} \right\} \left\{ \frac{z_1}{qw_1} \right\}}{\left\{ \frac{z_2}{qw_2} \right\} \left\{ \frac{qw_1}{z_1} \right\} \left\{ \frac{z_1}{qw_2} \right\} \left\{ \frac{qz_1}{w_1} \right\}} \Bigg) \\
& + (z_1 \leftrightarrow z_2) = 0.
\end{aligned} \tag{2.69}$$

We have used the abbreviations (2.41).

Proposition 2.7 The currents $\psi_j^\pm(z)$ ($1 \leq j \leq M + N - 1$) have the following formulae.

$$\psi_j^\pm(q^{\mp(r-\frac{c}{2})}z, p) = q^{\mp h_j} : \exp \left(- \sum_{m \neq 0} \frac{B_{j,m}}{[r^*m]_q} z^{-m} \right) : . \tag{2.70}$$

Here we have set

$$B_{j,m} = \begin{cases} \frac{[r^*m]_q}{[rm]_q} a_{j,m}, & (m > 0) \\ q^{c|m|} a_{j,m}, & (m < 0) \end{cases} \quad (1 \leq j \leq M + N - 1). \tag{2.71}$$

Definition 2.8 We define elliptic currents $E_j(z), F_j(z), H_j(z)$, ($1 \leq j \leq M + N - 1$) by

$$E_j(z) = e_j(z, p) e^{2Q_j} z^{-\frac{1}{r^*} P_j}, \tag{2.72}$$

$$F_j(z) = f_j(z, p) z^{\frac{1}{r} (P_j + h_j)}, \tag{2.73}$$

$$H_j^\pm(z) = H_j(q^{\pm(r-\frac{c}{2})}z), \tag{2.74}$$

$$H_j(z) =: \exp \left(- \sum_{m \neq 0} \frac{B_{j,m}}{[r^*m]_q} z^{-m} \right) : e^{2Q_j} z^{-\frac{c}{rr^*} P_j + \frac{1}{r} h_j}. \tag{2.75}$$

Here we have used the zero-mode operators P_j, Q_j , ($1 \leq j \leq M + N - 1$).

$$[P_i, Q_j] = -\frac{A_{i,j}}{2}, \quad (1 \leq i, j \leq M + N - 1). \quad (2.76)$$

Proposition 2.9 *The currents $E_j(z), F_j(z), H_j(z)$ and $B_{j,n}, h_j, c$, ($1 \leq j \leq M + N - 1, n \in \mathbb{Z}_{\neq 0}$) satisfy the defining relations of elliptic superalgebra $U_{q,p}(\widehat{sl}(M|N))$ (2.25), (2.26), (2.27), (2.28), (2.29), (2.30), (2.31), (2.32), (2.33), (2.34), (2.35), (2.36). They satisfy the Serre relations (2.37), (2.38) and (2.39), (2.40) for $1 \leq i, j \leq M + N - 1, (i \neq M)$ such that $|A_{i,j}| = 1$.*

We have constructed the elliptic deformed superalgebra $U_{q,p}(\widehat{sl}(M|N))$ from the quantum superalgebra $U_q(\widehat{sl}(M|N))$.

3 Bosonization

In this section we give new bosonization of the superalgebra $U_q(\widehat{sl}(1|2)), U_{q,p}(\widehat{sl}(1|2))$ for an arbitrary level k , and their screening currents. Wakimoto [41] constructed bosonization of affine algebra $\widehat{sl}(2)$ for an arbitrary level k . We call this-type bosonization based on the flag manifold [43] the Wakimoto realization. Feigin-Frenkel [42] generalized the Wakimoto realization to the higher-rank affine algebra $\widehat{sl}(N)$. Shiraishi [44] constructed the Wakimoto realization of the quantum algebra $U_q(\widehat{sl}(2))$ and its screening currents. Awata-Odake-Shiraishi constructed the Wakimoto realization for the quantum algebra $U_q(\widehat{sl}(N))$ and its screening currents [45]. In the case of $U_q(\widehat{sl}(2|1))$ Awata-Odake-Shiraishi [46] constructed the Wakimoto realization and Zhang-Gould [47] constructed the screening currents.

3.1 $U_q(\widehat{sl}(1|2)), U_{q,p}(\widehat{sl}(1|2))$, Screening

In this section we give new bosonizations of $U_q(\widehat{sl}(1|2)), U_{q,p}(\widehat{sl}(1|2))$ and their screening currents. In this section we assume the central element $c = k \neq 1$. The Cartan matrix $(A_{i,j})_{0 \leq i, j \leq 2}$ of $\widehat{sl}(1|2)$ is given by

$$(A_{i,j})_{0 \leq i, j \leq 2} = \begin{pmatrix} 0 & -1 & 1 \\ -1 & 0 & 1 \\ 1 & 1 & -2 \end{pmatrix}. \quad (3.1)$$

The Cartan matrix of the classical part $sl(1|2)$ is written by

$$(A_{i,j})_{1 \leq i, j \leq 2} = ((\nu_i + \nu_{i+1})\delta_{i,j} - \nu_i\delta_{i,j+1} - \nu_{i+1}\delta_{i+1,j})_{1 \leq i, j \leq 2},$$

where we have set $\nu_1 = +, \nu_2 = \nu_3 = -$. Let us introduce the bosons and the zero-mode operators a_m^j, Q_a^j ($m \in \mathbb{Z}, j = 1, 2$) $b_m^{i,j}, Q_b^{i,j}, c_m^{i,j}, Q_c^{i,j}$ ($m \in \mathbb{Z}, 1 \leq i < j \leq 3$) by

$$[a_m^i, a_n^j] = \frac{[(k-1)m]_q [A_{i,j}m]_q}{m} \delta_{m+n,0}, \quad [a_m^i, Q_a^j] = (k-1)A_{i,j} \delta_{m,0}, \quad (3.2)$$

$$[b_m^{i,j}, b_n^{i',j'}] = -\nu_i \nu_j \frac{[m]_q^2}{m} \delta_{i,i'} \delta_{j,j'} \delta_{m+n,0}, \quad [b_m^{i,j}, Q_b^{i',j'}] = -\nu_i \nu_j \delta_{i,i'} \delta_{j,j'} \delta_{m,0}, \quad (3.3)$$

$$[c_m^{i,j}, c_n^{i',j'}] = \nu_i \nu_j \frac{[m]_q^2}{m} \delta_{i,i'} \delta_{j,j'} \delta_{m+n,0}, \quad [c_m^{i,j}, Q_c^{i',j'}] = \nu_i \nu_j \delta_{i,i'} \delta_{j,j'} \delta_{m,0}. \quad (3.4)$$

Let us set the bosonic fields $a(z)$, $a_\pm(z)$ and $\left(\frac{1}{\beta} a\right)(z|\alpha)$ as follows.

$$a(z) = - \sum_{m \neq 0} \frac{a_m}{[m]_q} z^{-m} + Q_a + a_0 \log z, \quad (3.5)$$

$$a_\pm(z) = \pm(q - q^{-1}) \sum_{m > 0} a_{\pm m} z^{\mp m} \pm a_0 \log q, \quad (3.6)$$

$$\left(\frac{1}{\beta} a\right)(z|\alpha) = - \sum_{m \neq 0} \frac{a_m}{[\beta m]} q^{-\alpha|m|} z^{-m} + \frac{1}{\beta} (Q_a + a_0 \log z). \quad (3.7)$$

We impose the cocycle condition to the zero-mode operator.

$$e^{Q_b^{1,2}} e^{Q_b^{1,3}} = -e^{Q_b^{1,3}} e^{Q_b^{1,2}}, \quad e^{Q_b^{1,2}} e^{Q_b^{2,3}} = e^{Q_b^{2,3}} e^{Q_b^{1,2}}, \quad e^{Q_b^{1,2}} e^{Q_b^{2,3}} = e^{Q_b^{2,3}} e^{Q_b^{1,3}}. \quad (3.8)$$

Straightforward OPE calculations show the following propositions.

Proposition 3.1 *Bosonization of the quantum superalgebra $U_{q,p}(\widehat{sl}(1|2))$ is given as follows.*

$$c = k, \quad h_1 = a_0^1 - b_0^{2,3} - b_0^{1,2}, \quad h_2 = a_0^2 + 2b_0^{2,3} + b_0^{1,3} - b_0^{1,2}, \quad (3.9)$$

$$a_{1,m} = a_m^1 q^{-\frac{k-1}{2}|m|} - b_m^{2,3} q^{-(k-1)|m|} - b_m^{1,3} q^{-(k-1)|m|}, \quad (3.10)$$

$$a_{2,m} = a_m^2 q^{-\frac{k-1}{2}|m|} + b_m^{2,3} q^{-(k-1)|m|} (q^m + q^{-m}) + b_m^{1,3} q^{-(k-2)|m|} - b_m^{1,2} q^{-(k-1)|m|}, \quad (3.11)$$

$$x_1^+(z) = c_{1,1}^+ x_{1,1}^+(z) + c_{1,2}^+ x_{1,2}^+(z), \quad (3.12)$$

$$x_2^+(z) = \frac{1}{(q - q^{-1})z} (c_{2,1}^+ x_{2,1}^+(z) - c_{2,2}^+ x_{2,2}^+(z)), \quad (3.13)$$

$$x_1^-(z) = \frac{1}{(q - q^{-1})z} (c_{1,1}^- x_{1,1}^-(z) - c_{1,2}^- x_{1,2}^-(z) - c_{1,3}^- x_{1,3}^-(z) + c_{1,4}^- x_{1,4}^-(z)), \quad (3.14)$$

$$x_2^-(z) = \frac{1}{(q - q^{-1})z} (c_{2,1}^- x_{2,1}^-(z) - c_{2,2}^- x_{2,2}^-(z)) + c_{2,3}^- x_{2,3}^-(z), \quad (3.15)$$

where we have set

$$x_{1,1}^+(z) =: e^{-(b^{2,3} + b^{1,3})_+ (q^{-1}z) - b^{1,2} (q^{-1}z)} :, \quad (3.16)$$

$$x_{1,2}^+(z) =: e^{-(b+c)^{2,3}(z) - b^{1,3}(z)} :, \quad (3.17)$$

$$x_{2,1}^+(z) =: e^{b_+^{2,3}(z) + (b+c)^{2,3}(q^{-1}z)} :, \quad (3.18)$$

$$x_{2,2}^+(z) =: e^{b_{-}^{2,3}(z)+(b+c)^{2,3}(qz)} :, \quad (3.19)$$

$$x_{1,1}^-(z) =: e^{a_+^1(q^{\frac{k-1}{2}}z)+b^{1,2}(q^{k-1}z)} :, \quad (3.20)$$

$$x_{1,2}^-(z) =: e^{a_-^1(q^{-\frac{k-1}{2}}z)+b^{1,2}(q^{-k+1}z)} :, \quad (3.21)$$

$$x_{1,3}^-(z) =: e^{a_-^1(q^{-\frac{k-1}{2}}z)-b_-^{2,3}(q^{-k+1}z)+(b+c)^{2,3}(q^{-k}z)-b_-^{1,3}(q^{-k+1}z)+b^{1,3}(q^{-k}z)} :, \quad (3.22)$$

$$x_{1,4}^-(z) =: e^{a_-^1(q^{-\frac{k-1}{2}}z)-b_+^{2,3}(q^{-k+1}z)-b_-^{1,3}(q^{-k+1}z)+(b+c)^{2,3}(q^{-k+2}z)+b^{1,3}(q^{-k}z)} :, \quad (3.23)$$

$$x_{2,1}^-(z) =: e^{a_+^2(q^{\frac{k-1}{2}}z)+b_+^{2,3}(q^{k-2}z)-(b+c)^{2,3}(q^{k-1}z)+b_+^{1,3}(q^{k-2}z)-b_+^{1,2}(q^{k-1}z)} :, \quad (3.24)$$

$$x_{2,2}^-(z) =: e^{a_-^2(q^{-\frac{k-1}{2}}z)+b_-^{2,3}(q^{-k+2}z)-(b+c)^{2,3}(q^{-k+1}z)+b_-^{1,3}(q^{-k+2}z)-b_-^{1,2}(q^{-k+1}z)} :, \quad (3.25)$$

$$x_{2,3}^-(z) =: e^{a_+^2(q^{\frac{k-1}{2}}z)+b^{1,3}(q^{k-1}z)-b_+^{1,2}(q^{k-1}z)-b^{1,2}(q^{k-2}z)} :. \quad (3.26)$$

Here we have set the coefficients as follows.

$$(c_{1,1}^+, c_{1,2}^+, c_{2,1}^+, c_{2,2}^+) = (\alpha, \beta, \gamma, \gamma), \quad (3.27)$$

$$(c_{1,1}^-, c_{1,2}^-, c_{1,3}^-, c_{1,4}^-, c_{2,1}^-, c_{2,2}^-, c_{2,3}^-) = \left(\frac{1}{q\alpha}, \frac{1}{q\alpha}, \frac{1}{\beta}, \frac{1}{\beta}, \frac{1}{\gamma}, \frac{1}{\gamma}, \frac{q^{k-1}\alpha}{\beta\gamma} \right). \quad (3.28)$$

Here $\alpha, \beta, \gamma \neq 0$ are arbitrary parameters.

Next we give bosonization of the elliptic superalgebra $U_{q,p}(\widehat{sl}(1|2))$. Our construction is based on the dressing procedure of the quantum algebra developed in this paper.

Proposition 3.2 *Bosonization of the elliptic superalgebra $U_{q,p}(\widehat{sl}(1|2))$ is given as follows.*

$$c = k, \quad h_1 = a_0^1 - b_0^{2,3} - b_0^{1,2}, \quad h_2 = a_0^2 + 2b_0^{2,3} + b_0^{1,3} - b_0^{1,2}, \quad (3.29)$$

$$B_{j,m} = \begin{cases} \frac{[r^*m]_q}{[rm]_q} a_{j,m}, & (m > 0) \\ q^{k|m|} a_{j,m}, & (m < 0) \end{cases}, \quad (j = 1, 2), \quad (3.30)$$

$$a_{1,m} = a_m^1 q^{-\frac{k-1}{2}|m|} - b_m^{2,3} q^{-(k-1)|m|} - b_m^{1,3} q^{-(k-1)|m|}, \quad (3.31)$$

$$a_{2,m} = a_m^2 q^{-\frac{k-1}{2}|m|} + b_m^{2,3} q^{-(k-1)|m|} (q^m + q^{-m}) + b_m^{1,3} q^{-(k-2)|m|} - b_m^{1,2} q^{-(k-1)|m|}, \quad (3.32)$$

$$E_j(z) = u_j^+(z, p) x_j^+(z) e^{2Q_j} z^{-\frac{1}{r^*} P_j}, \quad (j = 1, 2), \quad (3.33)$$

$$F_j(z) = x_j^-(z) u_j^-(z, p) z^{\frac{1}{r} (P_j + h_j)}, \quad (j = 1, 2), \quad (3.34)$$

$$H_j^\pm(z) = H_j(q^{\pm(r-\frac{c}{2})}z), \quad (j = 1, 2), \quad (3.35)$$

where we have used (3.12), (3.13), (3.14), (3.15) and

$$u_j^+(z, p) = \exp \left(\sum_{m \geq 0} \frac{q^{r^*m}}{[r^*m]_q} B_{j,-m} z^m \right), \quad (j = 1, 2), \quad (3.36)$$

$$u_j^-(z, p) = \exp \left(- \sum_{m>0} \frac{q^{rm}}{[r^*m]_q} B_{j,m} z^{-m} \right), \quad (j = 1, 2), \quad (3.37)$$

$$H_j(z) =: \exp \left(- \sum_{m \neq 0} \frac{B_{j,m}}{[r^*m]_q} z^{-m} \right) : e^{2Q_j} z^{-\frac{c}{rr^*} P_j + \frac{1}{r} h_j}, \quad (j = 1, 2). \quad (3.38)$$

Here we have used the zero-mode operators

$$[P_i, Q_j] = -\frac{A_{i,j}}{2}, \quad (1 \leq i, j \leq 2). \quad (3.39)$$

Proposition 3.3 *The bosonic operators $s_j(z)$ ($j = 1, 2$) given below are the screening currents that commute with the quantum superalgebra $U_q(\widehat{sl}(1|2))$ modulo total difference.*

$$s_j(z) =: e^{-(\frac{1}{k-1}a^j)(z_1|\frac{k-1}{2})} \tilde{s}_j(z) : \quad (j = 1, 2). \quad (3.40)$$

Here we have set

$$\tilde{s}_1(z) = -c_{1,5} \tilde{s}_{1,5}(z), \quad (3.41)$$

$$\tilde{s}_2(z) = \frac{1}{(q - q^{-1})z} (-c_{2,3} \tilde{s}_{2,3}(z) + c_{2,4} \tilde{s}_{2,4}(z)) + c_{2,5} \tilde{s}_{2,5}(z), \quad (3.42)$$

where

$$\tilde{s}_{1,5}(z) = : e^{-b^{1,2}(z)} :, \quad (3.43)$$

$$\tilde{s}_{2,3}(z) = : e^{-b_+^{2,3}(qz) + (b+c)^{2,3}(q^2z) - b_-^{1,3}(qz) + b_-^{1,2}(z)} :, \quad (3.44)$$

$$\tilde{s}_{2,4}(z) = : e^{-b_-^{2,3}(qz) + (b+c)^{2,3}(z) - b_-^{1,3}(qz) + b_-^{1,2}(z)} :, \quad (3.45)$$

$$\tilde{s}_{2,5}(z) = : e^{-b^{1,3}(z) + b_+^{1,2}(z) + b^{1,2}(q^{-1}z)} :. \quad (3.46)$$

Here we have set the coefficients as follows.

$$(c_{1,5}, c_{2,3}, c_{2,4}, c_{2,5}) = \left(q\alpha, \gamma, \gamma, \frac{\beta\gamma}{q\alpha} \right), \quad (3.47)$$

where parameters $\alpha, \beta, \gamma \neq 0$ have been introduced in (3.27), (3.28) for the bosonizations of $U_q(\widehat{sl}(1|2))$. Explicitly the bosonic operators $s_1(z), s_2(z)$ and $x_1^\pm(z), x_2^\pm(z)$ satisfy the following relations.

$$[a_{i,m}, s_j(z_2)] = 0, \quad (3.48)$$

$$[x_i^+(z_1), s_j(z_2)] = 0, \quad (3.49)$$

$$\begin{aligned} [x_i^-(z_1), s_j(z_2)] &= \frac{\delta_{i,j}}{(q - q^{-1})z_1 z_2} (\delta(q^{k-1}z_2/z_1) - \delta(q^{-k+1}z_2/z_1)) \\ &\times : e^{-(\frac{1}{k-1}a^i)(z_1|\frac{k-1}{2})} :, \end{aligned} \quad (3.50)$$

and

$$\{\tilde{s}_1(z_1), \tilde{s}_1(z_2)\} = 0, \quad (3.51)$$

$$(z_1 - q^{-A_{1,2}} z_2) \tilde{s}_1(z_1) \tilde{s}_2(z_2) = (q^{-A_{2,1}} z_1 - z_2) \tilde{s}_2(z_2) \tilde{s}_1(z_1), \quad (3.52)$$

$$(z_1 - q^{-A_{2,2}} z_2) \tilde{s}_2(z_1) \tilde{s}_2(z_2) = (q^{-A_{2,2}} z_1 - z_2) \tilde{s}_2(z_2) \tilde{s}_2(z_1). \quad (3.53)$$

By the commutation relation $[a_{i,m}, s_j(z_2)] = 0$ we conclude the following.

Proposition 3.4 *The bosonic operators $s_j(z)$ ($j = 1, 2$) given in (3.40) become the screening currents that commute with the elliptic algebra $U_{q,p}(\widehat{sl}(1|2))$ modulo total difference. Explicitly the bosonic operators $s_1(z), s_2(z)$ and $E_1(z), E_2(z), F_1(z), F_2(z)$ satisfy the following relations.*

$$[B_{i,m}, s_j(z_2)] = 0, \quad (3.54)$$

$$[E_i(z_1), s_j(z_2)] = 0, \quad (3.55)$$

$$\begin{aligned} [F_i(z_1), s_j(z_2)] &= \frac{\delta_{i,j}}{(q - q^{-1})z_1 z_2} (\delta(q^{k-1} z_2/z_1) - \delta(q^{-k+1} z_2/z_1)) \\ &\times : e^{-(\frac{1}{k-1}a^i)(z_1| - \frac{k-1}{2})} u_i^-(z_1, p) z_1^{\frac{1}{r}(P_i + h_i)} : . \end{aligned} \quad (3.56)$$

The Jackson integral with parameters p and $s \neq 0$ is defined by

$$\int_0^{s\infty} f(z) d_p z = s(1-p) \sum_{n \in \mathbb{Z}} f(sp^n) p^n. \quad (3.57)$$

From the above proposition we have

$$\left[\int_0^{s\infty} s_j(z) d_{q^{2(k-1)}} z, U_{q,p}(\widehat{sl}(1|2)) \right] = 0. \quad (3.58)$$

3.2 $U_q(\widehat{sl}(2|1)), U_{q,p}(\widehat{sl}(2|1)),$ Screening

In this section we review known results on bosonization of $U_q(\widehat{sl}(2|1))$ [46] and its screening currents [47]. We give bosonizations of $U_{q,p}(\widehat{sl}(2|1))$ and its screenings. In this section we assume the central element $c = k \neq -1$. The Cartan matrix $(A_{i,j})_{0 \leq i,j \leq 2}$ of $\widehat{sl}(2|1)$ is given by

$$(A_{i,j})_{0 \leq i,j \leq 2} = \begin{pmatrix} 0 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 0 \end{pmatrix}. \quad (3.59)$$

The Cartan matrix of the classical part $sl(2|1)$ is written by

$$(A_{i,j})_{1 \leq i,j \leq 2} = ((\nu_i + \nu_{i+1})\delta_{i,j} - \nu_i\delta_{i,j+1} - \nu_{i+1}\delta_{i+1,j})_{1 \leq i,j \leq 2},$$

where we have set $\nu_1 = \nu_2 = +, \nu_3 = -$. Let us introduce the bosons and the zero-mode operators $a_m^j, Q_a^j, (m \in \mathbb{Z}, j = 1, 2)$ $b_m^{i,j}, Q_b^{i,j}, c_m^{i,j}, Q_c^{i,j} (m \in \mathbb{Z}, 1 \leq i < j \leq 3)$ by

$$[a_m^i, a_n^j] = \frac{[(k+1)m]_q [A_{i,j}m]_q}{m} \delta_{m+n,0}, \quad [a_m^i, Q_a^j] = (k+1)A_{i,j} \delta_{m,0}, \quad (3.60)$$

$$[b_m^{i,j}, b_n^{i',j'}] = -\nu_i \nu_j \frac{[m]_q^2}{m} \delta_{i,i'} \delta_{j,j'} \delta_{m+n,0}, \quad [b_m^{i,j}, Q_b^{i',j'}] = -\nu_i \nu_j \delta_{i,i'} \delta_{j,j'} \delta_{m,0}, \quad (3.61)$$

$$[c_m^{i,j}, c_n^{i',j'}] = \nu_i \nu_j \frac{[m]_q^2}{m} \delta_{i,i'} \delta_{j,j'} \delta_{m+n,0}, \quad [c_m^{i,j}, Q_c^{i',j'}] = \nu_i \nu_j \delta_{i,i'} \delta_{j,j'} \delta_{m,0}. \quad (3.62)$$

We impose the cocycle condition to the zero-mode operators.

$$e^{Q_b^{1,2}} e^{Q_b^{1,3}} = e^{Q_b^{1,3}} e^{Q_b^{1,2}}, \quad e^{Q_b^{1,2}} e^{Q_b^{2,3}} = e^{Q_b^{2,3}} e^{Q_b^{1,2}}, \quad e^{Q_b^{1,3}} e^{Q_b^{2,3}} = -e^{Q_b^{2,3}} e^{Q_b^{1,3}}. \quad (3.63)$$

Proposition 3.5 [46] *Bosonization of the quantum superalgebra $U_q(\widehat{sl}(2|1))$ is given as follows.*

$$c = k, \quad h_1 = a_0^1 + 2b_0^{1,2} + b_0^{1,3} - b_0^{2,3}, \quad h_2 = a_0^2 - b_0^{1,2} - b_0^{1,3}, \quad (3.64)$$

$$a_{1,m} = a_m^1 q^{-\frac{k+1}{2}|m|} + b_m^{1,2} q^{-(k+1)|m|} (q^m + q^{-m}) + b_m^{1,3} q^{-(k+2)|m|} - b_m^{2,3} q^{-(k+1)|m|}, \quad (3.65)$$

$$a_{2,m} = a_m^2 q^{-\frac{k+1}{2}|m|} - b_m^{1,2} q^{-(k+1)|m|} - b_m^{1,3} q^{-(k+1)|m|}, \quad (3.66)$$

$$x_1^+(z) = \frac{1}{(q - q^{-1})z} (c_{1,1}^+ x_{1,1}^+(z) - c_{1,2}^+ x_{1,2}^+(z)), \quad (3.67)$$

$$x_2^+(z) = c_{2,1}^+ x_{2,1}^+(z) + c_{2,2}^+ x_{2,2}^+(z), \quad (3.68)$$

$$x_1^-(z) = \frac{1}{(q - q^{-1})z} (c_{1,1}^- x_{1,1}^-(z) - c_{1,2}^- x_{1,2}^-(z)) + c_{1,3}^- x_{1,3}^-(z), \quad (3.69)$$

$$x_2^-(z) = \frac{1}{(q - q^{-1})z} (c_{2,1}^- x_{2,1}^-(z) - c_{2,2}^- x_{2,2}^-(z) - c_{2,3}^- x_{2,3}^-(z) + c_{2,4}^- x_{2,4}^-(z)), \quad (3.70)$$

where we have set

$$x_{1,1}^+(z) =: e^{b_+^{1,2}(z) - (b+c)^{1,2}(qz)}, \quad (3.71)$$

$$x_{1,2}^+(z) =: e^{b_-^{1,2}(z) - (b+c)^{1,2}(q^{-1}z)}, \quad (3.72)$$

$$x_{2,1}^+(z) =: e^{-b_+^{1,2}(qz) - b_+^{1,3}(qz) + b^{2,3}(qz)}, \quad (3.73)$$

$$x_{2,2}^+(z) =: e^{(b+c)^{1,2}(z) + b^{1,3}(z)}, \quad (3.74)$$

$$x_{1,1}^-(z) =: e^{a_+^1(q^{\frac{k+1}{2}}z) + b_+^{1,2}(q^{k+2}z) + (b+c)^{1,2}(q^{k+1}z) + b_+^{1,3}(q^{k+2}z) - b_+^{2,3}(q^{k+1}z)}, \quad (3.75)$$

$$x_{1,2}^-(z) =: e^{a_-^1(q^{-\frac{k+1}{2}}z) + b_-^{1,2}(q^{-k-2}z) + (b+c)^{1,2}(q^{-k-1}z) + b_-^{1,3}(q^{-k-2}z) - b_-^{2,3}(q^{-k-1}z)}, \quad (3.76)$$

$$x_{1,3}^-(z) =: e^{a_+^1(q^{\frac{k+1}{2}}z) - b_+^{2,3}(q^{k+1}z) - b^{1,3}(q^{k+1}z) + b^{2,3}(q^{k+1}z)}, \quad (3.77)$$

$$x_{2,1}^-(z) =: e^{a_+^2(q^{\frac{k+1}{2}}z) - b^{2,3}(q^{k+1}z)}, \quad (3.78)$$

$$x_{2,2}^-(z) =: e^{a_-^2(q^{-\frac{k+1}{2}}z) - b_{2,3}^{2,3}(q^{-k-1}z)} :, \quad (3.79)$$

$$x_{2,3}^-(z) =: e^{a_-^2(q^{-\frac{k+1}{2}}z) - b_{-}^{1,2}(q^{-k-1}z) - b_{-}^{1,3}(q^{-k-1}z) - (b+c)^{1,2}(q^{-k}z) - b^{1,3}(q^{-k}z)} :, \quad (3.80)$$

$$x_{2,4}^-(z) =: e^{a_-^2(q^{-\frac{k+1}{2}}z) - b_{+}^{1,2}(q^{-k-1}z) - b_{-}^{1,3}(q^{-k-1}z) - (b+c)^{1,2}(q^{-k-2}z) - b^{1,3}(q^{-k}z)} :. \quad (3.81)$$

Here we have set the coefficients as follows.

$$(c_{1,1}^+, c_{1,2}^+, c_{2,1}^+, c_{2,2}^+) = (\alpha, \alpha, \beta, \gamma), \quad (3.82)$$

$$(c_{1,1}^-, c_{1,2}^-, c_{1,3}^-, c_{2,1}^-, c_{2,2}^-, c_{2,3}^-, c_{2,4}^-) = \left(\frac{1}{\alpha}, \frac{1}{\alpha}, \frac{q^{k+1}\beta}{\alpha\gamma}, \frac{q}{\beta}, \frac{q}{\beta}, \frac{1}{\gamma}, \frac{1}{\gamma} \right). \quad (3.83)$$

Here $\alpha, \beta, \gamma \neq 0$ are arbitrary parameters.

Note. The coefficients of the currents $x_j^\pm(z)$ have 4 free parameters in [46]. In this paper we have only three free parameters α, β, γ , because we assume the commutation relations (3.102), (3.103), (3.104) with the screening currents.

Proposition 3.6 *Bosonization of the elliptic superalgebra $U_{q,p}(\widehat{sl}(2|1))$ is given as follows.*

$$c = k, \quad h_1 = a_0^1 + 2b_0^{1,2} + b_0^{1,3} - b_0^{2,3}, \quad h_2 = a_0^2 - b_0^{1,2} - b_0^{1,3}, \quad (3.84)$$

$$B_{j,m} = \begin{cases} \frac{[r^*m]_q}{[rm]_q} a_{j,m}, & (m > 0) \\ q^{k|m|} a_{j,m}, & (m < 0) \end{cases}, \quad (j = 1, 2), \quad (3.85)$$

$$a_{1,m} = a_m^1 q^{-\frac{k+1}{2}|m|} + b_m^{1,2} q^{-(k+1)|m|} (q^m + q^{-m}) + b_m^{1,3} q^{-(k+2)|m|} - b_m^{2,3} q^{-(k+1)|m|}, \quad (3.86)$$

$$a_{2,m} = a_m^2 q^{-\frac{k+1}{2}|m|} - b_m^{1,2} q^{-(k+1)|m|} - b_m^{1,3} q^{-(k+1)|m|}, \quad (3.87)$$

$$E_j(z) = u_j^+(z, p) x_j^+(z) e^{2Q_j} z^{-\frac{1}{r^*} P_j}, \quad (j = 1, 2), \quad (3.88)$$

$$F_j(z) = x_j^-(z) u_j^-(z, p) z^{\frac{1}{r} (P_j + h_j)}, \quad (j = 1, 2), \quad (3.89)$$

$$H_j^\pm(z) = H_j(q^{\pm(r-\frac{c}{2})} z), \quad (j = 1, 2), \quad (3.90)$$

where we have used (3.67), (3.68), (3.69), (3.70) and

$$u_j^+(z, p) = \exp \left(\sum_{m>0} \frac{q^{r^*m}}{[r^*m]_q} B_{j,-m} z^m \right), \quad (j = 1, 2), \quad (3.91)$$

$$u_j^-(z, p) = \exp \left(- \sum_{m>0} \frac{q^{rm}}{[r^*m]_q} B_{j,m} z^{-m} \right), \quad (j = 1, 2), \quad (3.92)$$

$$H_j(z) =: \exp \left(- \sum_{m \neq 0} \frac{B_{j,m}}{[r^*m]_q} z^{-m} \right) : e^{2Q_j} z^{-\frac{c}{rr^*} P_j + \frac{1}{r} h_j}, \quad (j = 1, 2). \quad (3.93)$$

Here we have used the zero-mode operators

$$[P_i, Q_j] = -\frac{A_{i,j}}{2}, \quad (1 \leq i, j \leq 2). \quad (3.94)$$

Proposition 3.7 [47] *The bosonic operators $s_1(z), s_2(z)$ given below are the screening currents that commute with the quantum superalgebra $U_q(\widehat{sl}(2|1))$ modulo total difference.*

$$s_j(z) = : e^{-(\frac{1}{k+1}a^j)(z|\frac{k+1}{2})} \tilde{s}_j(z) : \quad (j = 1, 2). \quad (3.95)$$

Here we have set

$$\tilde{s}_1(z) = \frac{1}{(q - q^{-1})z} (-c_{1,3}\tilde{s}_{1,3}(z) + c_{1,4}\tilde{s}_{1,4}(z)) + c_{1,5}\tilde{s}_{1,5}(z), \quad (3.96)$$

$$\tilde{s}_2(z) = -c_{2,5}\tilde{s}_{2,5}(z), \quad (3.97)$$

where

$$\tilde{s}_{1,5}(z) = : e^{b^{1,3}(z) - b^{2,3}(qz) + b_+^{2,3}(z)} :, \quad (3.98)$$

$$\tilde{s}_{1,4}(z) = : e^{-b_-^{1,2}(q^{-1}z) - (b+c)^{1,2}(z) + b_-^{2,3}(z) - b_-^{1,3}(q^{-1}z)} :, \quad (3.99)$$

$$\tilde{s}_{1,3}(z) = : e^{-b_+^{1,2}(q^{-1}z) - (b+c)^{1,2}(q^{-2}z) + b_-^{2,3}(z) - b_-^{1,3}(q^{-1}z)} :, \quad (3.100)$$

$$\tilde{s}_{2,5}(z) = : e^{b^{2,3}(z)} :.$$

Here we have set the coefficients as follows.

$$(c_{1,3}, c_{1,4}, c_{1,5}, c_{2,5}) = \left(\alpha, \alpha, \frac{q\alpha\beta}{\gamma}, \frac{\beta}{q} \right), \quad (3.101)$$

where parameters $\alpha, \beta, \gamma \neq 0$ have been introduced in (3.82), (3.83) for the bosonizations of $U_q(\widehat{sl}(2|1))$. Explicitly the bosonic operators $s_1(z), s_2(z)$ and $x_1^\pm(z), x_2^\pm(z)$ satisfy the following relations.

$$[a_{i,m}, s_j(z_2)] = 0, \quad (3.102)$$

$$[x_i^+(z_1), s_j(z_2)] = 0, \quad (3.103)$$

$$\begin{aligned} [x_i^-(z_1), s_j(z_2)] &= \frac{\delta_{i,j}}{(q - q^{-1})z_1 z_2} (\delta(q^{k+1}z_2/z_1) - \delta(q^{-k-1}z_2/z_1)) \\ &\times : e^{-(\frac{1}{k+1}a^i)(z_1|-\frac{k+1}{2})} :, \end{aligned} \quad (3.104)$$

and

$$(z_1 - q^{-A_{1,1}}z_2)\tilde{s}_1(z_1)\tilde{s}_1(z_2) = (q^{-A_{1,1}}z_1 - z_2)\tilde{s}_1(z_2)\tilde{s}_1(z_1), \quad (3.105)$$

$$(z_1 - q^{-A_{1,2}}z_2)\tilde{s}_1(z_1)\tilde{s}_2(z_2) = (q^{-A_{2,1}}z_1 - z_2)\tilde{s}_2(z_2)\tilde{s}_1(z_1), \quad (3.106)$$

$$\{\tilde{s}_2(z_1), \tilde{s}_2(z_2)\} = 0. \quad (3.107)$$

By the commutation relation $[a_{i,m}, s_j(z_2)] = 0$ we conclude the following.

Proposition 3.8 *The bosonic operators $s_j(z)$ ($j = 1, 2$) given in (3.95) become the screening currents that commute with the elliptic algebra $U_{q,p}(\widehat{sl}(2|1))$ modulo total difference. Explicitly the bosonic operators $s_1(z), s_2(z)$ and $E_1(z), E_2(z), F_1(z), F_2(z)$ satisfy the following relations.*

$$[B_{i,m}, s_j(z_2)] = 0, \quad (3.108)$$

$$[E_i(z_1), s_j(z_2)] = 0, \quad (3.109)$$

$$\begin{aligned} [F_i(z_1), s_j(z_2)] &= \frac{\delta_{i,j}}{(q - q^{-1})z_1 z_2} (\delta(q^{k+1} z_2/z_1) - \delta(q^{-k-1} z_2/z_1)) \\ &\times : e^{-(\frac{1}{k+1}a^i)(z_1| - \frac{k+1}{2})} u_i^-(z_1, p) z_1^{\frac{1}{r}(P_i + h_i)} : . \end{aligned} \quad (3.110)$$

From the above proposition we have

$$\left[\int_0^{s\infty} s_j(z) d_{q^{2(k-1)}} z, U_{q,p}(\widehat{sl}(2|1)) \right] = 0. \quad (3.111)$$

Acknowledgments

This work is supported by the Grant-in-Aid for Scientific Research **C** (21540228) from Japan Society for Promotion of Science. The author is grateful to Pascal Baseilhac and the colleagues in University of Tours for kind invitation and warm hospitality during his stay in Tours.

A Bosonization

In appendix we summarize relations of bosonization for $U_q(\widehat{sl}(1|2))$ and its screening currents relating to the delta-function $\delta(z) = \sum_{m \in \mathbb{Z}} z^m$.

$$\{x_{1,1}^+(z_1), x_{1,1}^-(z_2)\} \quad (A.1)$$

$$= \frac{q}{z_1} \delta(q^k z_2/z_1) e^{a_+^1(q^{\frac{k-1}{2}} z_2) - b_+^{2,3}(q^{k-1} z_2) - b_+^{1,3}(q^{k-1} z_2)},$$

$$\{x_{1,1}^+(z_1), x_{1,2}^-(z_2)\} \quad (A.2)$$

$$= \frac{q}{z_1} \delta(q^{-k+2} z_2/z_1) : e^{a_-^1(q^{-\frac{k-1}{2}} z_2) - b_+^{2,3}(q^{-k+2} z) - b_+^{1,3}(q^{-k+2} z_2)} :,$$

$$[x_{1,1}^+(z_1), x_{2,1}^-(z_2)] \quad (A.3)$$

$$= -(q - q^{-1}) \delta(q^{k-1} z_2/z_1) : e^{a_+^2(q^{\frac{k-1}{2}} z_2) - (b+c)^{2,3}(q^{k-1} z_2) - b_+^{1,2}(q^{k-1} z_2) - b^{1,2}(q^{k-2} z_2)} :,$$

$$\{x_{1,2}^+(z_1), x_{1,3}^-(z_2)\} \quad (\text{A.4})$$

$$= \frac{1}{z_1} \delta(q^{-k} z_2 / z_1) e^{a_-^1 (q^{-\frac{k-1}{2}} z_2) - b_-^{2,3} (q^{-k+1} z_2) - b_-^{1,3} (q^{-k+1} z_2)},$$

$$\{x_{1,2}^+(z_1), x_{1,4}^-(z_2)\} \quad (\text{A.5})$$

$$= \frac{1}{z_1} \delta(q^{-k+2} z_2 / z_1) : e^{a_-^1 (q^{-\frac{k-1}{2}} z_2) - b_+^{2,3} (q^{-k+2} z_2) - b_+^{1,3} (q^{-k+2} z_2)} :,$$

$$[x_{1,2}^+(z_1), x_{2,3}^-(z_2)] \quad (\text{A.6})$$

$$= -(q - q^{-1}) \delta(q^{k-1} z_2 / z_1) : e^{a_+^2 (q^{\frac{k-1}{2}} z_2) - (b+c)^{2,3} (q^{k-1} z_2) - b_+^{1,2} (q^{k-1} z_2) - b^{1,2} (q^{k-2} z_2)} :,$$

$$[x_{2,1}^+(z_1), x_{1,4}^-(z_2)] \quad (\text{A.7})$$

$$= -(q - q^{-1}) \delta(q^{-k+1} z_2 / z_1) : e^{a_-^1 (q^{-\frac{k-1}{2}} z_2) + (b+c)^{2,3} (q^{-k} z_2) + (b+c)^{2,3} (q^{-k+2} z_2) - b_-^{1,3} (q^{-k+1} z_2) + b^{1,3} (q^{-k} z_2)} :,$$

$$[x_{2,1}^+(z_1), x_{2,1}^-(z_2)] \quad (\text{A.8})$$

$$= (q - q^{-1}) \delta(q^k z_2 / z_1) e^{a_+^2 (q^{\frac{k-1}{2}} z_2) + b_+^{2,3} (q^{k-2} z_2) + b_+^{2,3} (q^k z_2) + b_+^{1,3} (q^{k-2} z_2) - b_+^{1,2} (q^{k-1} z_2)},$$

$$[x_{2,2}^+(z_1), x_{1,3}^-(z_2)] \quad (\text{A.9})$$

$$= (q - q^{-1}) \delta(q^{-k+1} z_2 / z_1) : e^{a_-^1 (q^{-\frac{k-1}{2}} z_2) + (b+c)^{2,3} (q^{-k} z_2) + (b+c)^{2,3} (q^{-k+2} z_2) - b_-^{1,3} (q^{-k+1} z_2) + b^{1,3} (q^{-k} z_2)} :,$$

$$[x_{2,2}^+(z_1), x_{1,3}^-(z_2)] \quad (\text{A.10})$$

$$= -(q - q^{-1}) \delta(q^{-k} z_2 / z_1) e^{a_-^2 (q^{-\frac{k-1}{2}} z_2) + b_-^{2,3} (q^{-k} z_2) + b_-^{2,3} (q^{-k+2} z_2) + b_-^{1,3} (q^{-k+2} z_2) - b_-^{1,2} (q^{-k+1} z_2)}.$$

$$[x_{2,1}^+(z_1), s_{2,3}(z_2)] \quad (\text{A.11})$$

$$= -(q - q^{-1}) \delta(q z_2 / z_1) : e^{-(\frac{1}{k-1} a^2) (z_2 | \frac{k-1}{2}) + b_-^{1,2} (z) - b_-^{1,3} (q z) + (b+c)^{2,3} (z) + (b+c)^{2,3} (q^2 z)} :,$$

$$[x_{2,2}^+(z_1), s_{2,4}(z_2)] \quad (\text{A.12})$$

$$= (q - q^{-1}) \delta(q z_2 / z_1) : e^{-(\frac{1}{k-1} a^2) (z_2 | \frac{k-1}{2}) + b_-^{1,2} (z) - b_-^{1,3} (q z) + (b+c)^{2,3} (z) + (b+c)^{2,3} (q^2 z)} :,$$

$$[x_{1,1}^+(z_1), s_{2,5}(z_2)] \quad (\text{A.13})$$

$$= \frac{1}{z_2} \delta(q^2 z_2 / z_1) : e^{-(\frac{1}{k-1} a^2) (z_2 | \frac{k-1}{2}) + b_-^{1,2} (z_2) - b_+^{1,3} (q z_2) - b^{1,3} (z_2) - b_+^{2,3} (q z_2)} :,$$

$$[x_{1,2}^+(z_1), s_{2,3}(z_2)] \quad (\text{A.14})$$

$$= q^{-1} (q - q^{-1}) \delta(q^2 z_2 / z_1) : e^{-(\frac{1}{k-1} a^2) (z_2 | \frac{k-1}{2}) + b_-^{1,2} (z_2) - b_+^{1,3} (q z_2) - b^{1,3} (z_2) - b_+^{2,3} (q z_2)} :,$$

$$\{x_{1,1}^-(z_1), s_{1,5}(z_2)\} = \frac{1}{z_2} \delta(q^{-k+1} z_2 / z_1) : e^{-(\frac{1}{k-1} a^1) (z_1 | -\frac{k-1}{2})} :, \quad (\text{A.16})$$

$$\{x_{1,2}^-(z_1), s_{1,5}(z_2)\} = \frac{1}{z_2} \delta(q^{k-1} z_2 / z_1) : e^{-(\frac{1}{k-1} a^1) (z_1 | -\frac{k-1}{2})} :, \quad (\text{A.17})$$

$$[x_{2,1}^-(z_1), s_{2,3}(z_2)] = (q - q^{-1}) \delta(q^{-k+3} z_2 / z_1)$$

$$\times : e^{a_+^2 (q^{\frac{k-1}{2}} z_1) - (\frac{1}{k-1} a^2) (q^{k-3} z_1 | \frac{k-1}{2}) + b_+^{1,3} (q^{k-2} z_1) - b_-^{1,3} (q^{k-2} z_1) - b_+^{1,2} (q^{k-1} z_1) + b_-^{1,2} (q^{k-3} z_1)} :, \quad (\text{A.18})$$

$$[x_{2,2}^-(z_1), s_{2,4}(z_2)] = -(q - q^{-1}) \delta(q^{k-1} z_2 / z_1) : e^{-(\frac{1}{k-1} a^2) (z_1 | -\frac{k-1}{2})} :, \quad (\text{A.19})$$

$$[x_{2,3}^-(z_1), s_{2,5}(z_2)] = \frac{-1}{(q - q^{-1}) q^{k-2} z_1 z_2}$$

$$\times (\delta(q^{-k+1}z_2/z_1) : e^{-(\frac{1}{k-1}a^2)(z_1|-\frac{k-1}{2})} : \quad (\text{A.20})$$

$$-\delta(q^{-k+3}z_2/z_1) : e^{a_+^2(q^{\frac{k-1}{2}}z_1)-(\frac{1}{k-1}a^2)(q^{k-3}z_1|\frac{k-1}{2})+b_+^{1,3}(q^{k-2}z_1)-b_-^{1,3}(q^{k-2}z_1)-b_+^{1,2}(q^{k-1}z_1)+b_-^{1,2}(q^{k-3}z_1)} : ,$$

$$[x_{1,2}^-(z_1), s_{2,3}(z_2)] \quad (\text{A.21})$$

$$= (q - q^{-1})\delta(q^k z_2/z_1) : e^{a_-^1(q^{\frac{k+1}{2}}z_2)-(\frac{1}{k-1}a^2)(z_2|\frac{k-1}{2})+b_-^{1,2}(z_2)+b^{1,2}(qz_2)-b_-^{1,3}(qz_2)-b_+^{2,3}(qz_2)+(b+c)^{2,3}(q^2z_2)} : ,$$

$$[x_{1,2}^-(z_1), s_{2,4}(z_2)] \quad (\text{A.22})$$

$$= (q - q^{-1})\delta(q^k z_2/z_1) : e^{a_-^1(q^{\frac{k+1}{2}}z_2)-(\frac{1}{k-1}a^2)(z_2|\frac{k-1}{2})+b_-^{1,2}(z_2)+b^{1,2}(qz_2)-b_-^{1,3}(qz_2)-b_-^{2,3}(qz_2)+(b+c)^{2,3}(q^2z_2)} : ,$$

$$[x_{1,3}^-(z_1), s_{2,3}(z_2)] = -q(q - q^{-1})\delta(q^k z_2/z_1) \quad (\text{A.23})$$

$$\times : e^{a_-^1(q^{\frac{k+1}{2}}z_2)-(\frac{1}{k-1}a^2)(z_2|\frac{k-1}{2})+b_-^{1,2}(z_2)-2b_-^{1,3}(qz_2)-b_-^{2,3}(qz_2)-b_+^{2,3}(qz_2)+(b+c)^{2,3}(z_2)+(b+c)^{2,3}(q^2z_2)} : ,$$

$$[x_{1,3}^-(z_1), s_{2,5}(z_2)] \quad (\text{A.24})$$

$$= \frac{1}{z_2}\delta(q^k z_2/z_1) : e^{a_-^1(q^{\frac{k+1}{2}}z_2)-(\frac{1}{k-1}a^2)(z_2|\frac{k-1}{2})+b_-^{1,2}(z_2)+b^{1,2}(qz_2)-b_-^{1,3}(qz_2)-b_-^{2,3}(qz_2)+(b+c)^{2,3}(q^2z_2)} : ,$$

$$[x_{1,4}^-(z_1), s_{2,4}(z_2)] = q(q - q^{-1})\delta(q^k z_2/z_1) \quad (\text{A.25})$$

$$\times : e^{a_-^1(q^{\frac{k+1}{2}}z_2)-(\frac{1}{k-1}a^2)(z_2|\frac{k-1}{2})+b_-^{1,2}(z_2)-2b_-^{1,3}(qz_2)-b_-^{2,3}(qz_2)-b_+^{2,3}(qz_2)+(b+c)^{2,3}(z_2)+(b+c)^{2,3}(q^2z_2)} : ,$$

$$[x_{1,4}^-(z_1), s_{2,5}(z_2)] \quad (\text{A.26})$$

$$= \frac{1}{z_2}\delta(q^k z_2/z_1) : e^{a_-^1(q^{\frac{k+1}{2}}z_2)-(\frac{1}{k-1}a^2)(z_2|\frac{k-1}{2})+b_-^{1,2}(z_2)+b^{1,2}(qz_2)-b_-^{1,3}(qz_2)-b_+^{2,3}(qz_2)+(b+c)^{2,3}(q^2z_2)} : .$$

References

- [1] A.A.Belavin, A.M.Polyakov and A.B.Zamolodchikov, Infinite Conformal Symmetry in Two-dimensional Quantum Field Theory, *Nucl.Phys.* **B241** 333-380 (1984).
- [2] M.Jimbo and T.Miwa, *Algebraic Analysis of Solvable Lattice Models*, CBMS Regional Conference Series in Mathematics **85** (American Mathematical Society), 1994.
- [3] G.E.Andrews, R.J.Baxter and P.J.Forrester, Eight-Vertex SOS Model and Generalized Rogers-Ramanujan-type identities, *J.Stat.Phys.* **35**, 193-266, (1984).
- [4] H.Konno, An Elliptic Algebra $U_{q,p}(\widehat{sl}_2)$ and the Fusion RSOS model, *Commun.Math.Phys.* **195**, 373-403, (1998).
- [5] M.Jimbo, H.Konno, S.Odake and J.Shiraishi, Elliptic Algebra $U_{q,p}(\widehat{sl}_2)$: Drinfeld Currents and Vertex Operators, *Commun.Math.Phys.* **199**, 605-647, (1999).

- [6] R.Baxter, Exactly Solved Models in Statistical Mechanics, Academic Press, London, 1982.
- [7] E.Date, M.Jimbo, T.Miwa and M.Okado, Fusion of the Eight Vertex SOS Model, *Lett.Math.Phys.***12**, 209-215, (1986).
- [8] E.Date, M.Jimbo, A.Kuniba, T.Miwa and M.Okado, Exactly Solvable SOS Models. II: Proof of the Star-triangle Relation and Combinatorial Identities, *Adv.Stu.Pure.Math.* **16**, 17-122, (1988).
- [9] E.Date, M.Jimbo, A.Kuniba, T.Miwa and M.Okado, Exactly Solvable SOS Models : Local Height Probabilities and Theta Function Identities, *Nucl.Phys.***B290**[FS20], 231-273, (1987).
- [10] M.Jimbo, T.Miwa and M.Okado, Solvable Lattice Models whose States are Dominant Integral Weights of $A_{n-1}^{(1)}$, *Lett.Math.Phys.***14**, 123-131, (1987).
- [11] M.Jimbo, A.Kuniba, T.Miwa and M.Okado, The $A_n^{(1)}$ Face Models, *Commun.Math.Phys.* **119**, 543-565, (1988).
- [12] B.Davies, O.Foda, M.Jimbo, T.Miwa and A.Nakayashiki, Diagonalization of the XXZ Hamiltonian by Vertex operators, *Commun.Math.Phys.***151**, 89-153, (1993).
- [13] M.Jimbo, K.Miki, T.Miwa and A.Nakayashiki, Correlation Functions of the XXZ Model for $\Delta < -1$, *Phys.Lett.***B168**, 256-263, (1992).
- [14] O.Foda, M.Jimbo, T.Miwa, K.Miki and A.Nakayashiki, Vertex Operators in Solvable Lattice Models, *J.Math.Phys.***35**, 13-46, (1994).
- [15] S.Lukyanov and Ya.Pugai, Multi-point Local Height Probabilities in the Integrable RSOS Model, *Nucl.Phys.***B473**, 631-658, (1996).
- [16] Y.Asai, M.Jimbo, T.Miwa and Ya.Pugai, Bosonizations of Vertex Operators for $A_{n-1}^{(1)}$ Face Model, *J.Phys.***A29**:Math.Gen., 6595-6616, (1996).
- [17] H.Furutsu, T.Kojima and Y.-H.Quano, Type-II Vertex Operators for the $A_{n-1}^{(1)}$ Face Model, *Int.J.Mod.Phys.***A15**, 1533-1556, (2000).
- [18] B.Feigin, M.Jimbo, T.Miwa, A.Odesskii and Ya.Pugai, Algebra of Screening Operators for the Deformed W_n Algebra, *Commun.Math.Phys.*—bf 191, 501-541, (1998).
- [19] M.Okado, Solvable Face Models Related to the Lie Superalgebra $sl(m|n)$, *Lett.Math.Phys.* **22**, 39-43, (1991).

- [20] V.G. Drinfeld, A New Realization of Yangians and Quantized Affine Algebras, *Sov.Math.Dokl.* **36**, 212-216, (1988).
- [21] T.Kojima and H.Konno, The Elliptic Algebra $U_{q,p}(\widehat{sl}_N)$ and the Drinfeld Realization of Elliptic Quantum Group $\mathcal{B}_{q,\lambda}(\widehat{sl}_N)$, *Commun.Math.Phys.* **239**, 405-447, (2003).
- [22] T.Kojima and H.Konno, The Drinfeld Realization of the Elliptic Quantum Group $\mathcal{B}_{q,\lambda}(A_2^{(2)})$, *J.Math.Phys.* **45**, 3146-3179, (2004).
- [23] E.K.Sklyanin, Some Algebraic Structure Connected with the Yang-Baxter Equation, *Funct.Anal.Appl.* **16**, 263-270, (1982).
- [24] O.Foda, K.Iohara, M.Jimbo, R.Kedem, T.Miwa and H.Yan, An Elliptic Quantum Algebra for \widehat{sl}_2 , *Lett.Math.Phys.* **32**, 259-268, (1994).
- [25] G.Felder, Elliptic Quantum Group, *Proc.ICMP Paris 1994*, Cambridge-Hong Kong : International Press, 211-218, (1995).
- [26] B.Enriquez and G.Felder, Elliptic Quantum Groups $E_{\tau,\eta}(\widehat{sl}_2)$ and Quasi-Hopf Algebras, *Commun.Math.Phys.* **195**, 651-689, (1998).
- [27] C.Frønsdal, Generalization and Exact Deformations of Quantum Groups, *Publ.Res.Math.Sci.* **33**, 91-149, (1997).
- [28] C.Frønsdal, Quasi-Hopf Deformation of Quantum Groups, *Lett.Math.Phys.* **40**, 117-134, (1997).
- [29] M.Jimbo, H.Konno, S.Odake and J.Shiraishi, Quasi-Hopf Twistors for Elliptic Quantum Groups, *Transformation Groups* **4**, 303-327, (1999).
- [30] Y.-Z. Zhang and M.D.Gould, Quasi-Hopf Superalgebras and Elliptic Quantum Superalgebra, *J.Math.Phys.* **40**, 5264-5282, (1999).
- [31] H.Konno, Elliptic quantum group $U_{q,p}(\widehat{sl}_2)$, Hopf algebroid structure and elliptic hypergeometric series, *J. Geometry and Physics* **59**, 1485-1511, (2009).
- [32] H.Konno, Elliptic quantum group and vertex operators, *J.Phys.* **A41**:Math.Theor. 194012(12 pages), (2008).
- [33] K.Miki, Creation/annihilation Operators and Form Factors of the XXZ model, *Phys.Lett.* **A186**, 217-224, (1994).

- [34] V.G.Kac and D.H.Peterson, Infinite-dimensional Lie Algebras, Theta Functions and Modular Forms, *Adv.Math.***53**, 125-264, (1984).
- [35] P.Goddard, A.Kent and D.Olive, Virasoro Algebras and Coset Space Models, *Phys.Lett.***B152**, 88-92, (1985).
- [36] V.Serganova, Kazhdan-Lusztig polynomials and character formula for the Lie superalgebra $gl(m|n)$, *Selecta Math.***2**, 607-651, (1996).
- [37] B.Jonathan, Kazhdan-Lusztig polynomials and character formula for the Lie superalgebra $gl(m|n)$, *J.Amer.Math.Sci.***16**, 185-231, (2003).
- [38] V.G.Kac, Lie Superalgebras, *Advances in Math.* **26**, 8-96, (1977).
- [39] Johan W.van de Leur, A Classification of Contragredient Lie Superalgebras of Finite Growth, *Commun. in Algebra* **17**, 1815-1841, (1989).
- [40] H.Yamane, On Defining Relations of the affine Lie Superalgebras and their Quantized Universal Enveloping Superalgebras, *Publ.Res.Inst.Math.Sci.* **35**, 321-390, (1999).
- [41] M.Wakimoto, Fock Representation of the Affine Lie Algebra $A_1^{(1)}$, *Commun.Math.Phys.* **104**, 605-609, (1986).
- [42] B.L.Feigin and E.V.Frenkel, Representation of Affine Kac-Moody Algebras and Bosonization, *Physics and Mathematics of Strings* (World Scientific), 271-316, 1990.
- [43] B.L.Feigin and E.V.Frenkel, Affine Kac-Moody Algebras and Semi-Infinite Flag Manifolds, *Commun.Math.Phys.* **128**, 161-189, (1990).
- [44] J.Shiraishi, Free Boson Representation of $U_q(\widehat{sl}_2)$, *Phys.Lett.* **A171**, 243-248, (1992).
- [45] H.Awata, S.Odake and J.Shiraishi, Free Boson Realization of $U_q(\widehat{sl}_N)$, *Commun.Math.Phys.* **162**, 61-83, (1994).
- [46] H.Awata, S.Odake and J.Shiraishi, q -Difference Realization of $U_q(sl(M|N))$ and Its Application to Free Boson Realization of $U_q(\widehat{sl}(2|1))$, *Lett.Math.Phys.* **42**, 271-279, (1997).
- [47] Y.-Z.Zhang and M.D.Gould, $U_q(\widehat{sl}(2|1))$ Vertex Operators, Screen Currents and Correlation Functions at Arbitrary Level, *J.Math.Phys.***41**, 5277-5291, (2000).